

Protein kinase C isoforms: Multi-functional regulators of cell life and death

Mary E. Reyland

Department of Craniofacial Biology, University of Colorado Health Sciences Center, Aurora, CO 80045, USA

TABLE OF CONTENTS

1. Abstract
2. Introduction
 - 2.1. The PKC superfamily
 - 2.2. Modes of regulation
3. PKC isoforms that regulate cell survival
 - 3.1. PKC-alpha
 - 3.2. PKC-epsilon
 - 3.3. PKC-zeta and PKC-lambda/iota
4. PKC isoforms that regulate apoptosis
 - 4.1. Regulation of apoptosis by PKC-delta
 - 4.2. Activation of PKC-delta by apoptotic signals
 - 4.2.1. Tyrosine phosphorylation of PKC-delta
 - 4.2.2. Nuclear localization of PKC-delta.
 - 4.2.3. Caspase-cleavage of PKC-delta
5. PKC and disease: A potential role in human cancer
6. Conclusions
7. References

1. ABSTRACT

The protein kinase C (PKC) family consists of 10 related serine/threonine protein kinases some of which are critical regulators of cell proliferation, survival and cell death. While early studies relied on broad spectrum chemical activators or inhibitors of this family, the generation of isoform specific tools has greatly facilitated our understanding of the contribution of specific PKC isoforms to cell proliferation and apoptosis. These studies suggest that PKC-alpha, PKC-epsilon, and the atypical PKC's, PKC-lambda/iota and PKC-zeta, preferentially function to promote cell proliferation and survival, while the novel isoform, PKC-delta is an important regulator of apoptosis. The essential role of this kinase family in both cell survival and apoptosis suggests that specific isoforms may function as molecular sensors, promoting cell survival or cell death depending on environmental cues. Given their central role in cell and tissue homeostasis, it is not surprising that the expression or activity of some of these kinases is altered in human diseases, particularly cancer.

2. INTRODUCTION

The PKC family consists of 10 serine/threonine protein kinases originally characterized by their dependency upon lipids for catalytic activity (1-4). The past 30 years of investigation has revealed roles for PKC in the regulation of a plethora of cellular processes. Early studies relied on activation of PKC by phorbol-12-myristate-13-acetate (PMA) and/or inhibition by pharmacological agents to demonstrate a role for PKC in the regulation of specific cell functions. More recently, the complexity and potential redundancy of the PKC signaling network has prompted the development of PKC isoform specific tools including dominant inhibitory kinases, mouse models in which specific PKC isoforms are disrupted, and PKC isoform specific antisense/siRNA to define isoform-specific functions of PKC. These tools have provided evidence that PKC isoforms regulate a variety of essential biological processes including cell migration, contraction, immunity, neural plasticity, proliferation, differentiation, apoptosis and metabolism (5). A remarkable finding is that the




Figure 1. The PKC superfamily. PKC isoforms contain four structurally conserved domains (C1-C4). The N-terminal regulatory domain contains a pseudosubstrate binding site (PS), and the DAG (C1) and Ca^{++} (C2, or “C2-like”) binding sites. The catalytic terminal domain contains the ATP binding domain (C3), the substrate binding site, and the kinase domain (C4). The regulatory and catalytic domains are separated by a flexible hinge region which is the site of cleavage of PKC-delta by caspase-3 in apoptotic cells.

function of a specific PKC isoform can vary between different cell types, implying that the specification of responses may rely on interaction of PKC isoforms with anchoring proteins or other regulatory pathways in the cell. This review will focus on the biology of the PKC isoform family and discuss the role of specific isoforms in the regulation of cell proliferation and apoptosis. The implications of alterations in the activation/expression of this important kinase family in human cancer will also be considered.

2.1. The PKC superfamily

Protein kinase C was discovered by Yasutomi Nishizuka in 1977 when his group purified a cyclic nucleotide independent, Ca^{2+} and lipid-dependent, kinase from rat and bovine cerebellum (1, 2). Southern hybridization using low stringency probes later revealed a family of related isoforms (6). In mammals, the PKC superfamily is divided into conventional, novel and atypical subfamilies based on isoform structure and requirement for specific lipid and ion co-activators (3, 5) [see Figure 1]. PKC's are also found in lower eukaryotes; *C. elegans* has four PKC's that represent all classes of mammalian isoforms, while *S. cerevisiae* has a single PKC (Pkc1) that resembles the mammalian atypical PKCs (7). The remarkable conservation of this kinase family from yeast to humans suggests that PKC plays a critical role in cellular signaling, and analysis of their role in simple systems has provided valuable insights into more complex mammalian systems.

2.2. Modes of regulation

The PKC superfamily, together with cAMP-dependent kinase (PKA) and cGMP-dependent kinase (PKG), comprises a related family of enzymes, the “AGC” kinases (8). The AGC kinases consist of a catalytic core whose activity is allosterically controlled by a regulatory domain. The lipid dependence of the PKC enzyme family has greatly facilitated the identification of upstream activators of these kinases. Physiologic regulators of PKC,

including growth factors and hormones, induce the generation of diacylglycerol (DAG) and the release of intracellular Ca^{++} . Sub-families of PKC are defined by their requirement for these activators. The conventional isoforms (PKC-alpha, -beta I, -beta II and -gamma) require DAG and Ca^{++} , the novel isoforms (PKC-delta, -epsilon, -eta and -theta) require DAG, but not Ca^{++} , while the atypical isoforms (PKC-zeta and lambda/iota) require neither.

The conventional and novel PKC isoforms are structurally conserved, consisting of a N-terminal regulatory domain which contains a pseudosubstrate binding site, and the DAG and Ca^{++} cofactor binding C1 and C2 domains, respectively (8). The C-terminal domain is highly conserved between isoforms and contains the ATP binding domain, the substrate binding site, and the catalytic domain (Figure 1). Phosphorylation at three sites in C-terminal domain is required to generate a mature form of the kinase that can be activated by lipids (9-15). It should be appreciated that these are constitutive, and not stimulus regulated, phosphorylation events and thus are not thought to play a role in determining the specificity of isoform activity. The first of these phosphorylation events occurs at a conserved threonine in the activation loop; phosphorylation at this site appears to be essential for activity of most isoforms (12, 15). Several laboratories have identified the PIP3 regulated kinase, PDK-1, as the kinase responsible for PKC activation loop phosphorylation (10, 16). Phosphorylation at two additional C-terminal sites contributes to the stability of the kinase. These include an autophosphorylation site and a C-terminal hydrophobic site (12). Phosphorylation at these sites renders PKC protease and phosphatase resistant, and catalytically competent; however phosphorylated PKC is still in an inactive conformation in which the substrate binding pocket is occupied by the pseudosubstrate domain.

It is thought that most cellular PKC is fully phosphorylated and in an auto-inhibited conformation in the cytoplasm. Stimulus induced generation of DAG and Ca^{++} results in allosteric activation of the conventional and novel PKC isoforms via a number of well defined steps (4, 12). First, binding of DAG to the C1 domain increases the affinity of PKC for the membrane. In the case of isoforms that are also regulated by Ca^{++} , Ca^{++} is thought to increase the affinity of the C1 domain for DAG. Second, interaction with acidic membrane lipids provides the energy necessary to dislodge the pseudosubstrate from the substrate binding pocket and activate the kinase. Activated PKC can then bind and phosphorylate substrates at the membrane. Notably, while membrane translocation is often used as a read out of PKC activation, “active” PKC is known to be targeted to other locations within the cell. This targeting may follow activation of the kinase at the membrane, or PKC may be activated at non-membrane sites perhaps by lipid-independent mechanisms (see discussion below). For instance, the PKC N-terminal regulatory domain contains binding sites for anchoring proteins; interaction with these anchoring proteins is hypothesized to target the activated kinase to specific sub-cellular sites and/or substrates. These include the

Receptors for Activated C Kinase (RACK's) as well as other PKC-interacting proteins (17, 18). Interaction of PKC with anchoring proteins may help to explain the substrate specificity of PKC isoforms, a perplexing question since multiple isoforms can be activated downstream of a given signal.

While lipid-dependent conventional and novel isoforms are thought to be largely activated by the "canonical" pathway described above, little is known about how the lipid-independent atypical PKC's (PKC-zeta and PKC-lambda/iota) are activated. Recent studies suggest that the activity of these isoforms may be regulated through protein-protein interactions involving the PB-1 domain of the kinase. Interaction of PKC-lambda/iota with Prostate Apoptosis Response-6 (PAR-6), and PKC-zeta with Prostate Apoptosis Response-4 (PAR-4), both require the PB-1 domain of the kinase and these interactions have been shown to modulate kinase activity (19, 20). Notably, candidate drugs that disrupt interaction of the PB-1 domain of PKC-lambda with PAR-6 have been identified and shown to inhibit the transformed growth of lung cancer cell lines (19). It is likely that in response to some stimuli conventional and novel PKC isoforms may also be activated by alternative, lipid-independent pathways. This includes allosteric regulation via protein-protein interactions, tyrosine phosphorylation, and perhaps most dramatically, caspase cleavage of PKC-delta in apoptotic cells to generate a constitutively active kinase (21, 22). A common theme of these alternative activation mechanisms appears to be modification of interactions between the catalytic and regulatory domain, or in the case of caspase cleavage, removal of the regulatory domain, to allow exposure of the substrate binding site and possibly other C-terminal motifs.

3. PKC ISOFORMS THAT REGULATE CELL SURVIVAL

The tumor promoting properties of phorbol esters have been known for many years and well documented in animal models of human cancer. The discovery of PKC as the phorbol ester "receptor" lead to a heightened interest in the contribution of these kinases to tumorigenesis and tumor progression. In fact, as discussed below, altered expression of many PKC isoforms has been documented in human cancer, suggesting that these family members may function as oncogenes or tumor suppressors. Here we will discuss those PKC isoforms that are associated primarily with a pro-survival phenotype. However it should be appreciated that given the impact of cellular context on phenotype, attempts at categorizing specific PKC isoforms as pro-survival or pro-apoptotic will most certainly be inaccurate at times.

3.1. PKC-alpha

Numerous studies suggest that PKC-alpha promotes cell survival and that loss of PKC-alpha activity sensitizes cells to apoptotic agents. Whether PKC-alpha enhances cell survival by promoting cell proliferation is controversial. On one hand, overexpression of PKC-alpha increases proliferation in many cell types, including mouse

fibroblasts, MCF-7 breast cancer cells and U87 malignant glioma cells, and depletion of PKC-alpha reduces the proliferation of lung cancer cells (23-26). On the other hand, some studies, including *in vivo* tumor studies (see below), suggest that PKC-alpha can negatively regulate cell proliferation and that loss of PKC-alpha function results in increased proliferation (27, 28).

Both the positive and negative effects of PKC-alpha on cell proliferation appear to be mediated via effects on the cell cycle machinery. For instance, PKC-alpha promotes cell cycle progression by increasing transcription of cyclin D1 in 3T3 fibroblasts and through up regulation of the CDK inhibitor, p21, in glioma cells (25, 29). However, PKC-alpha can also suppress proliferation in hepatoma cells, intestinal epithelial cells and pancreatic cancer cells through down regulation of cyclin D1 and/or up regulation of the CDK inhibitors, p21 and p27 (30-34). Intestinal epithelial cells have provided a useful model to investigate the signaling pathways downstream of PKC-alpha in suppression of proliferation. Studies in these cells suggest that PKC-alpha -dependent activation of the Ras/Raf/MAPK and/or Akt pathways may account for many of the effects of PKC-alpha on cell cycle regulators, including cyclin D1, p21 and p27 (31, 35). PKC-alpha can suppress cyclin D1 translation through Akt-dependent activation of the translational repressor, 4E-BP1 (31). PKC-alpha can also regulate the Akt pathway via direct phosphorylation of Akt at serine 473, and through activation of the serine/threonine protein kinase Raf-1 (36-39).

There is abundant evidence that PKC-alpha may support survival by suppressing apoptosis. In glioma cells, salivary epithelial cells, and melanoma and bladder carcinoma cell lines, depletion or inhibition of PKC-alpha induces apoptosis (23, 40-42). Likewise, Heregulin-induced apoptosis is enhanced by down regulation of PKC-alpha (43). Our studies in salivary epithelial cells indicate that the absence of PKC-alpha induces a PKC-delta-dependent apoptotic program (40). However, treatment of LNCaP prostate cancer cells with a synthetic DAG analog that activates PKC-alpha, but not PKC-delta, was shown to induce apoptosis (44, 45). Apoptosis induced by loss of PKC-alpha maybe secondary to loss of a proliferative signal, or may occur through a direct effect on the apoptotic machinery. In support of the later possibility, proteins involved in the execution of apoptosis have been identified as targets of PKC-alpha. Overexpression of PKC-alpha suppresses apoptosis in human pre-B REH cells and increases Bcl-2 phosphorylation at serine 70 which stabilizes and increases the anti-apoptotic function of Bcl-2 (46-48). Conversely, depletion of PKC-alpha in COS cells induces apoptosis and down regulation of Bcl-2 expression (42).

How can the discordant effects of PKC-alpha on proliferation, cell cycle progression and apoptosis be reconciled? Studies indicate that these divergent outcomes may be mediated at least in part by PKC-alpha-induced increases in the expression of p21. Although known primarily as a cell cycle inhibitor, p21 can also promote

cell cycle progression and inhibit apoptosis, presumably through its interaction with distinct regulatory complexes (49, 50). A recent study by Oliva et al indicates that activation of PKC- α can also induce senescence of non-small cell lung cancer cells through increased expression of p21 (51). In non-dividing cells, PKC- α mediated increases in p21 may contribute to cell cycle inhibition and maintenance of the senescent state. However in the context of a proliferative cell, p21 may drive proliferation through interaction with a set of cell cycle regulators distinct from those that regulate senescence. The dual role of PKC- α in promoting and suppressing the cell cycle suggests that the cellular context may play a significant role in determining the outcome of PKC- α activation.

3.2. PKC- ϵ

PKC- ϵ plays a well established role in cell proliferation (52). Overexpression of PKC- ϵ in NIH 3T3 cells or Rat 6 fibroblasts results in increased growth rate, loss of contact inhibition and tumors in nude mice (53). Furthermore, overexpression and activation of PKC- ϵ in mouse epidermis results in squamous cell carcinomas, consistent with its proposed oncogenic function (54, 55). Downstream effectors of PKC- ϵ associated with proliferation and cell transformation include the Ras/Raf/MAPK pathway and Signal Transducer and Activator of Transcription 3 (STAT 3) (54, 55). Activation of the Ras/Raf/MAPK pathway is necessary for transformation by PKC- ϵ , as expression of a dominant negative Raf-1 can reverse the PKC- ϵ mediated transformation of Rat 6 fibroblasts and colonic epithelial cells (52, 56-58).

Suppression of apoptosis may contribute to PKC- ϵ -mediated tumorigenesis (59-63). Early *in vitro* studies showed that PKC- ϵ is required for PMA mediated protection of U937 cells from TNF- α or calphostin C induced apoptosis (64). Subsequent studies have demonstrated an anti-apoptotic role for PKC- ϵ in most, but not all, cell models investigated. In some cases PKC- ϵ may function down stream of growth factors to suppress apoptosis. In endothelial cells, activation of PKC- ϵ downstream of VEGF can suppress apoptosis associated with vascular injury (62). In small cell lung cancer cells, PKC- ϵ is required for suppression of apoptosis downstream of FGF-2 activation (61). Likewise, overexpression of PKC- ϵ generally protects against Fas and TRAIL-induced cell death (59, 60, 65). Caspase cleavage of PKC- ϵ also occurs in some cells undergoing apoptosis; in glioma cells caspase cleavage of PKC- ϵ contributes to the TRAIL-induced apoptotic program (66-68).

PKC- ϵ also protects against apoptosis induced by DNA damaging agents, and has been implicated in the resistance of tumor cells to anti-cancer drugs. Elevated PKC- ϵ levels are observed in chemoresistance non-small cell lung cancer cell lines and in chemo-resistant ovarian cells (63, 69). Consistent with these observations, overexpression of PKC- ϵ can promote survival of lung cancer cells and increase their resistance to chemotherapeutic drugs (63). In breast cancer cells PKC- ϵ controls Akt activation, and this

appears to contribute to the suppression of apoptosis observed in breast cancer cells in which PKC- ϵ is overexpressed (59). PKC- ϵ may also directly inhibit apoptosis through up regulation of the anti-apoptotic protein, Bcl-2, or downregulation of pro-apoptotic proteins such as Bid (60, 62).

3.3. PKC- λ /iota and PKC- ζ

The atypical PKC isoforms, PKC-iota (and the mouse homolog, PKC- λ) and PKC- ζ are critical for cell survival signaling, presumably due to their role as downstream effectors of PI-3 kinase (70). Most of the effects of PKC-iota and PKC- ζ can be attributed to activation of the pro-survival NF- κ B signaling pathway and suppression of apoptosis (20, 71, 72). Many studies indicate a correlation between the expression or activation of PKC-iota and/or PKC- ζ and sensitivity to apoptosis. PKC- ζ suppresses Fas-induced apoptosis in Jurkat cells, and PKC- ζ inhibition increases apoptosis in leukemia cells exposed to etoposide and TNF- α (73, 74). Conversely, overexpression of PKC- ζ suppresses apoptosis induced by chemotherapeutic agents in U937 cells; topoisomerase II was identified as a downstream target of PKC- ζ in these studies (75). Due to the high homology between the PKC-iota and PKC- ζ isoforms, tools to distinguish their individual functions have been limited. However recent studies in knock-out mice indicate distinct roles for these isoforms, as PKC-iota, but not PKC- ζ , is critical for mouse development (20). Studies by Murray and Fields suggest that PKC-iota may be the more important of these isoforms in the context of suppression of apoptosis. Their studies show that PKC-iota, but not PKC- ζ , protects leukemia cells from chemotherapeutic agents and that Bcr-Abl mediated resistance to apoptosis is mediated by PKC-iota (19, 76, 77).

One of the more interesting aspects of the atypical PKC's is their dynamic regulation by agents which induce cell death. In some contexts, transient activation of these isoforms is observed in response to apoptotic agents. PKC-iota/PKC- ζ are activated by caspase cleavage which results in its degradation through the ubiquitin-proteasome pathway (78, 79). Likewise, atypical PKC expression is increased prior to cell death induced by UV, ceramide, and in cytokine-mediated cartilage destruction (80-82). PKC- ζ activity in apoptotic cells is negatively regulated by its interaction with the pro-apoptotic protein, PAR-4, which suppresses NF- κ B activation (71, 83-85). Sequestration of PKC- ζ and PKC- λ by PAR-4 is one mechanism by which PAR-4 induces apoptosis. PAR-4 can also directly interact with and suppress the enzymatic activity of PKC- ζ and PKC- λ in apoptotic cells (86). PKC- ζ may also be inhibited by interaction with p38, as seen in apoptotic chondrocytes (87). Recently it has been shown that p38 interacts with the regulatory domain of PKC- ζ and suppresses PKC- ζ activity by blocking autophosphorylation of the kinase (88). Conversely, PKC- ζ and PKC- λ have been shown to bind to, and inactivate, the pro-survival kinase, Akt, in response to ceramide and growth factors, and to phosphorylate and inhibit the pro-apoptotic protein, Bad

(89-92). The cumulative data suggests that through regulation of pro-survival pathways, the atypical PKC's may act as a switch between cell survival and cell death.

4. PKC ISOFORMS THAT REGULATE APOPTOSIS

Both PKC-delta and PKC-theta can be cleaved by caspase-3 to generate a constitutively activated kinase, which, when introduced into cells, can induce apoptosis (93, 94). Based on this observation these kinases are often grouped together as "pro-apoptotic". However, there is only limited evidence that PKC-theta plays a significant role in apoptosis. PKC-theta appears instead to be critical for immune cell function and T cell survival, and PKC-theta null mice display severe defects in T-cell signaling and activation (95). In contrast, the function of PKC-delta in apoptotic cells has been well established.

PKC-delta is a ubiquitously expressed isoform and studies in PKC-delta-/- mice have identified diverse roles for this signaling molecule in control of proliferation, immunity, apoptosis and cell migration (96). While PKC-delta is clearly an important regulator of apoptosis, it also has been shown to suppress cell cycle progression and negatively regulate proliferation. Studies from Santiago-Walker, et al, suggest that the anti-proliferative effects of PKC-delta may contribute to induction of apoptosis (97). In these studies, the overexpression of PKC-delta stimulated progression through the G1 phase of the cell cycle, but induced S phase arrest followed by apoptosis (97). In contrast, some studies, particularly in transformed cells, support a role for PKC-delta in promoting cell cycle progression and proliferation (98). In this context PKC-delta may act downstream of growth factors such as IGF-1 to promote proliferation (99, 100). Here we will discuss data that supports a role for PKC-delta as an early regulator of apoptosis, and mechanisms which regulate pro-apoptotic signal transduction by this ubiquitously expressed kinase.

4.1. Regulation of apoptosis by PKC-delta

PKC-delta is activated by numerous apoptotic stimuli and is required for apoptosis induced by genotoxins (22, 101), oxidative stress (102) and death receptors (103). Treatment with the PKC-delta selective inhibitor, rottlerin (104), expression of kinase inactive PKC-delta (PKC-delta KD) (101) or the introduction of a PKC-delta specific RACK inhibitory peptide (105), have all been shown to inhibit apoptosis. We have used inhibitors of PKC-delta to probe the function of PKC-delta in the apoptotic pathway. Our studies show that expression of PKC-delta KD inhibits both downstream apoptotic events such as caspase activation and DNA fragmentation, as well as upstream apoptotic events such as loss of mitochondrial membrane potential (101, 104). These studies from our lab, together with numerous studies from other labs, point to an essential role for PKC-delta as a regulator of early events in the apoptotic pathway (101, 104, 106, 107). In support of this conclusion, smooth muscle cells and primary salivary epithelial cells derived from PKC-delta -/- mice are defective in mitochondria-dependent apoptosis (107, 108). Analysis of primary parotid cells from PKC-delta -/- mice shows a decrease in cytochrome c release, PARP cleavage

and caspase-3 activation in response to etoposide, however, apoptosis can be completely restored by reintroduction of wild type PKC-delta (107). Loss of PKC-delta also protects parotid glands against γ -irradiation induced apoptosis *in vivo* (107). Although apoptosis is suppressed, activation of p53 is normal in primary parotid cells from wild type and PKC-delta -/- mice, suggesting that PKC-delta is activated downstream of cellular damage signals (107).

Substrates of PKC-delta identified in apoptotic cells include transcription factors, protein kinases, structural proteins, DNA repair and checkpoint molecules and Bcl-2 family members. The majority of PKC-delta's substrates in apoptotic cells are nuclear proteins. PKC-delta interacts with and phosphorylates DNA-dependent protein kinase (DNA-PK) in cells exposed to genotoxins (109, 110). Phosphorylation of DNA-PK inhibits DNA binding, suggesting that PKC-delta mediated phosphorylation inactivates the DNA double strand break repair function of this protein (109). PKC-delta can also phosphorylate the checkpoint protein Rad9; phosphorylated Rad9 can then bind and sequester Bcl-2, inducing cell death (110). Other reports suggest that PKC-delta may directly target and inactivate the apoptosis machinery. PKC-delta has been shown to phosphorylate and activate capsase-3 (111), to phosphorylate and target the anti-apoptotic protein Mcl-1 for degradation (112), and to suppress phosphorylation of the pro-apoptotic protein, Bad (113). Finally, PKC-delta may regulate transcription through activation of p53. In one study, downregulation of PKC-delta inhibited basal transcription of p53, while other studies report PKC-delta dependent accumulation of the p53 protein in apoptotic cells (114-116).

PKC-delta has been shown to interface with downstream signaling cascades to regulate the apoptotic machinery. Our laboratory has shown that PKC-delta and Signal Transducer and Activator of Transcription 1 (STAT1) can interact in etoposide treated cells and that STAT1 is a downstream target of PKC-delta in the apoptotic pathway (115). In addition, there is accumulating evidence that the PI3-kinase/AKT pathway, the ERK, JNK and p38 pathways may be downstream targets of PKC-delta in apoptotic cells. In cardiomyocytes exposed to ischemia, Akt is dephosphorylated and inactivated in a PKC-delta dependent manner. Inactivation of Akt results in dephosphorylation and release of the pro-apoptotic protein, Bad, and initiation of cell death (113). PKC-delta enhances radiation-induced apoptosis via ERK1/2 activation and suppression of radiation-induced G2-M arrest. PKC-delta activates the JNK pathway in irradiation and Ara-c induced apoptosis, possibly through phosphorylation and activation of MEKK1 (117). Primary cells derived from PKC-delta -/- mice are deficient in JNK activation, and JNK activation *in vivo* in response to x-irradiation is reduced in these mice (107). In contrast, binding of HSP25 to PKC-delta inhibits cell death, suggesting that the pro-apoptotic function of PKC-delta may be negatively regulated through interaction with other proteins (118).




Figure 2. Regulation of the proapoptotic function of PKC-delta. Under normal growth conditions PKC-delta is retained in the cytoplasm by a mechanism that is dependent upon the regulatory domain. Apoptotic signals result in tyrosine phosphorylation of the regulatory domain and allow nuclear accumulation of PKC-delta. Active caspase-3 likewise accumulates in the nucleus resulting in cleavage of PKC-delta and generation of delta-CF. Delta-CF localizes constitutively in the nucleus where it may regulate proteins involved in the cell damage response.

4.2. Activation of PKC-delta by apoptotic signals

PKC-delta is ubiquitously expressed and regulates diverse cell functions, hence, under normal conditions the proapoptotic function of PKC-delta must be tightly regulated in order to assure cell survival. We have defined multiple events that regulate the pro-apoptotic function of PKC-delta and have shown that these events are temporally coordinated in apoptotic cells (see Figure 2). The first event appears to be transduction of a “death” signal to PKC-delta by a damage sensing pathway. Our studies suggest that this occurs through phosphorylation of PKC-delta on specific tyrosine residues (119). Activated PKC-delta then transiently accumulates in the nucleus (120). If caspase-3 is also translocated to the nucleus, nuclear PKC-delta is cleaved by caspase to generate the PKC-delta catalytic fragment (delta-CF), a constitutively activated, pro-apoptotic form of PKC-delta that accumulates in the nucleus (120). Nuclear accumulation of PKC-delta appears to be both necessary and sufficient to commit a cell to apoptosis, whereas cytoplasmic retention of PKC-delta in the cytoplasm assures cell survival (120).

4.2.1. Tyrosine phosphorylation of PKC-delta.

Studies from our lab and others suggest that phosphorylation of PKC-delta on tyrosine may be a mechanism to dictate stimulus specific functions of PKC-delta (119, 121, 122). Functionally important tyrosine residues in the context of apoptosis include in PKC-delta include Y64 and Y187 in glioma cells treated with etoposide (121) and Y311, Y332 and Y512 in response to H₂O₂ (122). We have shown that tyrosine phosphorylation of PKC-delta in response to apoptotic agents occurs very rapidly and that tyrosines 64 and 155 in the regulatory

domain must be phosphorylated for nuclear translocation of PKC-delta to occur (119). Non-receptor tyrosine kinases, specifically c-Abl, Src and Lyn have been implicated in the phosphorylation of PKC-delta in response to apoptotic agents (95, 123, 124).

4.2.2. Nuclear localization of PKC-delta

Many PKC-delta substrates in apoptotic cells are nuclear proteins, consistent with the nuclear translocation of PKC-delta in response to many apoptotic agents (125, 126). We have defined a nuclear localization sequence in the catalytic domain of PKC-delta and have shown that nuclear localization is required for regulation of apoptosis by PKC-delta (125). Our studies suggest that once tyrosine phosphorylated PKC-delta rapidly accumulates in the nucleus (119). Hence, tyrosine phosphorylation in the regulatory domain may facilitate the binding of nuclear import receptors to the PKC-delta nuclear localization signal. This suggests that the regulatory domain of PKC-delta functions in part to retain PKC-delta in the cytoplasm in the absence of an apoptotic signal.

4.2.3. Caspase-cleavage of PKC-delta

Cleavage of PKC-delta by caspase occurs in the hinge domain of the protein between the regulatory and catalytic domains and results in release of a constitutively active catalytic fragment (delta-CF). We have shown that cleavage of PKC-delta by caspase occurs in the nucleus of apoptotic cells and is dependent on the coordinated import of activated caspase-3 into the nucleus (120). Importantly, while PKC-delta transiently accumulates in the nucleus in response to genotoxins, delta-CF when generated is largely or entirely nuclear at the early stages of apoptosis (120). Thus caspase cleavage facilitates the sustained nuclear accumulation of PKC-delta, which appears to be critical for commitment to apoptosis in some cells (120). The delta-CF may have additional functions in apoptotic cells, particularly in the later stages of apoptosis when it is also found in the cytoplasm (120). For instance, Sitailo et al have shown that expression of delta-CF results in activation of the pro-apoptotic protein, Bax, and cytochrome c release in keratinocytes (127). Taken together, our studies suggest that regulating the subcellular locale of PKC-delta is critical for cell survival, and that caspase cleavage of PKC-delta, which allows its nuclear accumulation, signals an irreversible commitment to apoptosis.

5. PKC AND DISEASE: A POTENTIAL ROLE IN HUMAN CANCER

The ability of phorbol esters to promote tumors has been known for many years and well documented in animal models of human cancer. The discovery of PKC as the major phorbol ester “receptor” in the cell has lead to intensive investigation into the potential contribution of these kinases to human cancer (128, 129). Not surprisingly, the PKC isoforms most commonly associated with increased proliferation and/or survival, PKC-alpha and PKC-epsilon, are those most commonly overexpressed in human cancer, and represent potential oncogenes. To this group should be added PKC-iota, as increased expression of PKC-iota has recently been shown to correlate with

tumor stage in non-small cell lung cancer, and depletion of PKC- ϵ was shown to reverse the transformed phenotype of these cells (30). In contrast, the observation that loss of PKC- δ is associated with cell transformation in some cells, together with the preponderance of data suggesting an anti-proliferative/pro-apoptotic role for PKC- δ , has led to the suggestion the PKC- δ may function as a tumor suppressor (98, 130). Although this hypothesis has not been tested *in vivo*, PKC- δ expression is reduced in human squamous cell carcinomas, and PKC- δ expression decreases with increasing tumor grade in human endometrial carcinomas (131, 132).

Overexpression of PKC- α is seen in a variety of human tumors, suggesting that it may contribute to tumor progression (133-135). In particular, in human breast cancer cells PKC- α has been shown to regulate HER2/neu expression and contribute to HER2/neu-mediated cell invasion (43, 133). PKC- α antisense oligonucleotides have been shown in some instances to decrease proliferation and sensitize cells to apoptosis, and PKC- α antisense oligonucleotides reduce tumor growth in mouse xenographs (136-138). However, clinical trials using this approach for the treatment of human tumors using have been disappointing (139, 140). Given the overlapping roles of PKC- α in both promoting and suppressing cell cycle progression, its functional analysis *in vivo* is likely to be complex. In this regard, recent studies using mouse models in which PKC- α is inhibited or genetically “knocked-out” indicate that, perhaps contrary to expectation, that the loss of PKC- α can accelerate colon cancer and result in increased B cell proliferation and leukemia (27, 28).

Clinical studies suggest that increased PKC- ϵ expression is strongly associated with tumor progression in many human cancers. PKC- ϵ has been documented as a biomarker of aggressive breast cancer, and high PKC- ϵ expression correlates with poor prognosis and positive Her2/neu status (141). The PKC- ϵ gene is also amplified in 28% of thyroid cancers and a chimeric/truncated version of PKC- ϵ has been cloned from human thyroid cancer cells (142). Likewise, PKC- ϵ expression is increased in gliomas, and depletion of PKC- ϵ can trigger apoptosis in glioma cell lines possibly through an effect on Akt mediated survival (65, 143). Overexpression of PKC- ϵ is commonly seen in human prostate tumors, and is associated with conversion from an androgen dependent to androgen independent state (144, 145). PKC- ϵ may also contribute to the resistance of prostate cancer cells to apoptosis through interaction with the pro-apoptotic protein, Bax (146). Interestingly, increased association of PKC- ϵ with Bax correlates with the increased resistance to apoptosis in prostate cancer cells (146). In addition to its pro-proliferative properties, PKC- ϵ may regulate the invasion and metastasis of cancer cells (147).

With few exceptions, proof of a definitive role for specific PKC isoforms in cancer cell growth, invasion and survival has been elusive. This presumably reflects

both the paucity of *in vivo* models available, as well as the multiple functions many of these kinases appear to have *in vivo*. However, there is compelling evidence that PKC isoform expressions is altered in human tumors, suggesting that a subset of these kinases may function as tumor suppressors or oncogenes, or cooperate with other genes to promote human tumors (148).

6. CONCLUSIONS

Abundant evidence implicates PKC isoforms as critical regulators of cell proliferation and apoptosis. PKC kinases regulate cell survival and apoptosis both “upstream” via growth factor signaling, as well as “downstream” via regulation of transcription factors and/or posttranslational modification of proteins. As might be expected, alterations in the expression and/or activity of specific members of this family are associated with the pathogenesis of some human diseases. In most cases, the contribution of PKC isoforms to human disease appears to be related to their role as regulators of cell proliferation and apoptosis. Thus, understanding the molecular mechanisms by which specific isoforms regulate cellular signaling pathways may enable the identification of new therapeutic targets to treat human disease.

7. REFERENCES

1. Inoue, M., A. Kishimoto, Y. Takai & Y. Nishizuka: Studies on a cyclic nucleotide-independent protein kinase and its proenzyme in mammalian tissues. II. Proenzyme and its activation by calcium-dependent protease from rat brain. *J Biol Chem*, 252, 7610-7616 (1977)
2. Takai, Y., A. Kishimoto, M. Inoue & Y. Nishizuka: Studies on a cyclic nucleotide-independent protein kinase and its proenzyme in mammalian tissues. I. Purification and characterization of an active enzyme from bovine cerebellum. *J Biol Chem*, 252, 7603-7609 (1977)
3. Newton, A.: Regulation of the ABC kinases by phosphorylation: protein kinase C as a paradigm. *Biochemical J*, 370, 361-371 (2003)
4. Newton, A. C.: Protein kinase C: structural and spatial regulation by phosphorylation, cofactors, and macromolecular interactions. *Chem Rev*, 101, 2352-64 (2001)
5. Dempsey, E. C., A. C. Newton, D. Mochly-Rosen, A. P. Fields, M. E. Reyland, P. A. Insel & R. O. Messing: Protein kinase C isozymes and the regulation of diverse cell responses. *Am J Physiol*, 279, L429-438 (2000)
6. Ono, Y., T. Fujii, K. Ogita, U. Kikkawa, K. Igarashi & Y. Nishizuka: The structure, expression, and properties of additional members of the protein kinase C family. *J Biol Chem*, 263, 6927-6932 (1988)
7. Mellor, H. & P. J. Parker: The extended protein kinase C superfamily. *Biochem J*, 332, 281-292 (1998)

8. Gold, M. G., D. Barford & D. Komander: Lining the pockets of kinases and phosphatases. *Curr Opin Struct Biol*, 16, 693-701 (2006)
9. Bornancin, F. & P. J. Parker: Phosphorylation of protein kinase C- α on serine 657 controls the accumulation of active enzyme and contributes to its phosphatase-resistant state. *J Biol Chem*, 272, 3544-9 (1997)
10. Le Good, J. A., W. H. Ziegler, D. B. Parekh, D. R. Alessi, P. Cohen & P. J. Parker: Protein kinase C isotypes controlled by phosphoinositide 3-Kinase through the protein kinase PDK1. *Science*, 281, 2042-2045 (1998)
11. Parekh, D., W. Ziegler, K. Yonezawa, K. Hara & P. J. Parker: Mammalian TOR controls one of two kinase pathways acting upon nPKC- δ and nPKC- ϵ . *J Biol Chem*, 274, 34758-34764 (1999)
12. Parekh, D. B., W. Ziegler & P. J. Parker: Multiple pathways control protein kinase C phosphorylation. *EMBO J*, 19, 496-503 (2000)
13. Li, W., J. Zhang, D. P. Bottaro, W. Li & J. H. Pierce: Identification of serine 643 of protein kinase C- δ as an important autophosphorylation site for its enzymatic activity. *J Biol Chem*, 272, 24550-24555 (1997)
14. Stempka, L., M. Schnolzer, S. Radke, G. Rincke, F. Marks & M. Gschwendt: Requirements of protein kinase C- δ for catalytic function. Role of glutamic acid 500 and autophosphorylation of serine 643. *J Biol Chem*, 274, 8886-8892 (1999)
15. Stempka, L., A. Girod, H.-J. Muller, G. Rincke, F. Marks, M. Gschwendt & D. Bossemeyer: Phosphorylation of protein kinase C- δ at threonine 505 is not a prerequisite for enzymatic activity. Expression of rat PKC- δ and an alanine 505 mutant in bacteria in a functional form. *J Biol Chem*, 272, 6805-6811 (1997)
16. Dutil, E. M., A. Toker & A. Newton: Regulation of conventional protein kinase C isozymes by phosphoinositide-dependent kinase 1 (PDK-1) *Curr Biol*, 8, 1366-1375 (1998)
17. Jaken, S. & P. Parker: Protein kinase C binding partners. *BioEssays*, 22, 245-254 (2000)
18. Poole, A. W., G. Pula, I. Hers, D. Crosby & M. L. Jones: PKC-interacting proteins: from function to pharmacology. *Trends Pharm Sci*, 25, 528-535 (2004)
19. Fields, A. P. & R. P. Regala: Protein kinase C α : Human oncogene, prognostic marker and therapeutic target. *Pharmacol Res*, 55, 487-497 (2007)
20. Moscat, J., P. Rennert & M. T. Diaz-Meco: PKC ζ at the crossroad of NF- κ B and Jak1/Stat6 signaling pathways. *Cell Death Differ*, 13, 702-711 (2005)
21. Steinberg, S.: Distinctive activation mechanisms and functions for protein kinase C- δ . *Biochem J*, 384, 449-459 (2004)
22. Ghayur, T., M. Hugunin, R. V. Talanian, S. Ratnofsky, C. Quinlan, Y. Emoto, P. Pandey, R. Datta, Y. Huang, S. Kharbanda, H. Allen, R. Kamen, W. Wong & D. Kufe: Proteolytic activation of protein kinase C- δ by an ICE/CED 3-like protease induces characteristics of apoptosis. *J Exp Med*, 184, 2399-2404 (1996)
23. Mandil, R., E. Ashkenazi, M. Blass, I. Kronfeld, G. Kazimirsky, G. Rosenthal, F. Umansky, P. S. Lorenzo, P. M. Blumberg & C. Brodie: Protein kinase C- α and protein kinase C- δ play opposite roles in the proliferation and apoptosis of glioma cells. *Cancer Res*, 61, 4612-4619 (2001)
24. Ways, D. K., C. A. Kukoly, J. deVente, J. L. Hooker, W. O. Bryant, K. J. Posekany, D. J. Fletcher, P. P. Cook & P. J. Parker: MCF-7 breast cancer cells transfected with protein kinase C- α exhibit altered expression of other protein kinase C isoforms and display a more aggressive neoplastic phenotype. *J Clin Invest*, 95, 1906-1915 (1995)
25. Soh, J.-W. & I. B. Weinstein: Roles of specific isoforms of protein kinase C in the transcriptional control of cyclin D1 and related genes. *J Biol Chem*, 278, 34709-34716 (2003)
26. Yin, J., J. Gao, R. Shao, W. Tian, J. Wang & Y. Wan: siRNA agents inhibit oncogene expression and attenuate human tumor cell growth. *J Exp Ther Oncol*, 3, 194-204 (2003)
27. Nakagawa, R., J. W. Soh & A. M. Michie: Subversion of Protein kinase C α signaling in hematopoietic progenitor cells results in the generation of a B-cell chronic lymphocytic leukemia-like population *in vivo*. *Cancer Res*, 66, 527-534 (2006)
28. Oster, H. & M. Leitges: Protein Kinase C α but not PKC- ζ suppresses intestinal tumor formation in ApcMin/+ Mice. *Cancer Res*, 66, 6955-6963 (2006)
29. Besson, A. & V. W. Yong: Involvement of p21 waf1/Cip1 in protein kinase- α induced cell cycle progression. *Mol Cell Biol*, 20, 4580-4590 (2000)
30. Hizli, A. A., A. R. Black, M. A. Pysz & J. D. Black: Protein kinase C- α signaling inhibits cyclin D1 translation in intestinal epithelial cells. *J Biol Chem*, 281, 14596-14603 (2006)
31. Guan, L., K. Song, M. A. Pysz, K. J. Curry, A. A. Hizli, D. Danielpour, A. R. Black & J. D. Black: Protein kinase C-mediated down-regulation of cyclin D1 involves activation of the translational repressor 4E-BP1 via a phosphoinositide 3-kinase/Akt-independent, protein phosphatase 2A-dependent mechanism in intestinal epithelial cells. *J Biol Chem*, 282, 14213-14225 (2007)

32. Wu, T., Y.-H. Hsieh, Y.-S. Hsieh & J. Liu: Reduction of PKC- α decreases cell proliferation, migration, and invasion of human malignant hepatocellular carcinoma. *J Cell Biochem*, 103, 9-20 (2008)
33. Frey, M. R., J. A. Clark, O. Leontieva, J. M. Uronis, A. R. Black & J. D. Black: Protein kinase C signaling mediates a program of cell cycle withdrawal in the intestinal epithelium. *J Cell Biol*, 151, 763-778 (2000)
34. Detjen, K., F. Brembeck, M. Welzel, A. Kaiser, H. Haller, B. Wiedenmann & S. Rosewicz: Activation of protein kinase C α inhibits growth of pancreatic cancer cells via p21 (cip)-mediated G (1) arrest. *J Cell Sci*, 113, 3025-3035 (2000)
35. Clark, J. A., A. R. Black, O. V. Leontieva, M. R. Frey, M. A. Pysz, L. Kunneva, A. Woloszynska-Read, D. Roy & J. D. Black: Involvement of the ERK signaling cascade in protein kinase C-mediated cell cycle arrest in intestinal epithelial cells. *J Biol Chem*, 279, 9233-9247 (2004)
36. Kolch, W., G. Heidecker, G. Kocks, R. Hummel, H. Vahldt, N. Mischak, G. Finkenzeller, D. Marne & U. R. Rapp: Protein kinase C α activates RAF-1 by direct phosphorylation. *Nature*, 364, 249-252 (1993)
37. Li, L., K. Sampat, N. Hu, J. Zakari & S. H. Yuspa: Protein kinase C negatively regulates Akt activity and modifies UVC-induced apoptosis in mouse keratinocytes. *J Biol Chem*, 281, 3237-3243 (2006)
38. Partovian, C. & M. Simons: Regulation of protein kinase B/Akt activity and Ser473 phosphorylation by protein kinase C- α in endothelial cells. *Cell Signal*, 16, 951-957 (2004)
39. Li, W., J. Zhang, L. Flechner, T. Hyun, A. Yam, T. Franke & J. Pierce: Protein kinase C- α overexpression stimulates Akt activity and suppresses apoptosis induced by interleukin 3 withdrawal. *Oncogene*, 18, 6564-6572 (1999)
40. Matassa, A. A., R. L. Kalkofen, L. Carpenter, T. J. Biden & M. E. Reyland: Inhibition of PKC- α induces a PKC- δ -dependent apoptotic program in salivary epithelial cells. *Cell Death Differ*, 10, 269-277 (2003)
41. Jorgensen, K., M. Skrede, V. Cruciani, S.-O. Mikalsen, A. Slipicevic & V. A. Florenes: Phorbol ester phorbol-12-myristate-13-acetate promotes anchorage-independent growth and survival of melanomas through MEK-independent activation of ERK1/2. *Biochem Biophys Res Commun*, 329, 266-274 (2005)
42. Whelan, D. H. R. & P. J. Parker: Loss of protein kinase C function induces an apoptotic response. *Oncogene*, 16, 1939-1944 (1998)
43. Le, X., M. Marcelli, A. McWatters, B. Nan, G. Mills, C. O'Brian & J. Bast RC: Heregulin-induced apoptosis is mediated by down-regulation of Bcl-2 and activation of caspase-7 and is potentiated by impairment of protein kinase C- α activity. *Oncogene*, 20, 8258-8269 (2001)
44. Garcia-Bermejo, M. L., F. C. Leskow, T. Fujii, Q. Wang, P. M. Blumberg, M. Ohba, T. Kuroki, K. C. Han, J. Lee, V. E. Marquez & M. G. Kazanietz: Diacylglycerol (DAG)-lactones, a new class of protein kinase C (PKC) agonists, induce apoptosis in LNCaP prostate cancer cells by selective activation of PKC- α . *J Biol Chem*, 277, 645-55 (2002)
45. Gavrielides, M., A. Frijhoff, C. Conti & K. MG: Protein kinase C and prostate carcinogenesis: targeting the cell cycle and apoptotic mechanisms. *Curr Drug Targets*, 5, 431-443 (2004)
46. Ruvolo, P. P., X. Deng, B. K. Carr & W. S. May: A functional role for mitochondrial protein kinase C- α Bcl-2 phosphorylation and suppression of apoptosis. *J Biol Chem*, 273, 25436-25442 (1998)
47. Jiffar, T., S. Kurinna, G. Suck, D. Carlson-Bremer, M. R. Ricciardi, M. Konopleva, M. Andreeff & P. P. Ruvolo: PKC- α mediates chemoresistance in acute lymphoblastic leukemia through effects on Bcl2 phosphorylation. *Leukemia*, 18, 505-512 (2004)
48. Deng, X., T. Ito, B. Carr, M. Mumby & W. S. May Jr.: Reversible phosphorylation of Bcl2 following interleukin 3 or bryostatin 1 is mediated by direct interaction with protein phosphatase 2A. *J Biol Chem*, 273, 34157-34163 (1998)
49. Gartel, A. L. & A. L. Tyner: The role of the cyclin-dependent kinase inhibitor p21 in apoptosis. *Mol Cancer Ther*, 1, 639-649 (2002)
50. Gartel, A. L. & S. K. Radhakrishnan: Lost in transcription: p21 repression, mechanisms, and consequences. *Cancer Res*, 65, 3980-3985 (2005)
51. Oliva, J. L., M. C. Caino, A. M. Senderowicz & M. G. Kazanietz: S-phase-specific activation of PKC- α induces senescence in non-small cell lung cancer cells. *J Biol Chem*, 283, 5466-76 (2008)
52. Cacace, A. M., M. Ueffing, A. Philipp, E. K. Han, W. Kolch & I. B. Weinstein: PKC- ϵ functions as an oncogene by enhancing activation of the Raf kinase. *Oncogene*, 13, 2517-2626 (1996)
53. Mischak, H., J. A. Goodnight, W. Kolch, G. Martiny-Baron, C. Schaehtle, M. G. Kazanietz, P. M. Blumberg, J. H. Pierce & J. F. Mushinski: Overexpression of protein kinase C- δ and - ϵ in NIH 3T3 cells induces opposite effects on growth, morphology, anchorage dependence, and tumorigenicity. *J Biol Chem*, 268, 6090-6096 (1993)
54. Aziz, M. H., H. T. Manoharan, D. R. Church, N. E. Dreckschmidt, W. Zhong, T. D. Oberley, G. Wilding & A. K. Verma: Protein kinase C ϵ interacts with signal

transducers and activators of transcription 3 (Stat3), phosphorylates Stat3 Ser727, and regulates its constitutive activation in prostate cancer. *Cancer Res*, 67, 8828-8838 (2007)

55. Aziz, M. H., H. Moammir, H. Manoharan, J. Sand & A. Verma: Protein kinase C epsilon interacts with Stat3 and regulates its activation that is essential for the development of skin cancer. *Mol Carcinog*, 46, 646-653 (2007)

56. Perletti, G. P., M. Folini, H. C. Lin, H. Mischak, F. Piccinini & A. H. J. Tashjian: Overexpression of protein kinase C-epsilon is oncogenic in rat colonic epithelial cells. *Oncogene*, 12, 847-854 (1996)

57. Perletti, G. P., P. Concari, S. Brusaferrì, E. Marras, F. Piccinini & A. H. J. Tashjian: Protein kinase C-epsilon is oncogenic in colon epithelial cells by interaction with the ras signal transduction pathway. *Oncogene*, 16, 3345-3348 (1998)

58. Piiper, A., R. Elez, S.-J. You, B. Kronenberger, S. Loitsch, S. Roche & S. Zeuzem: Cholecystokinin stimulates extracellular signal-regulated kinase through activation of the epidermal growth factor receptor, Yes, and protein kinase C. SIGNAL AMPLIFICATION AT THE LEVEL OF Raf BY ACTIVATION OF PROTEIN KINASE C-epsilon. *J Biol Chem*, 278, 7065-7072 (2003)

59. Lu, D., J. Huang & A. Basu: Protein kinase C epsilon activates protein kinase B/Akt via DNA-PK to protect against tumor necrosis factor-alpha-induced cell death. *J Biol Chem*, 281, 22799-22807 (2006)

60. Sivaprasad, U., E. Shankar & A. Basu: Downregulation of Bid is associated with PKC-epsilon mediated TRAIL resistance. *Cell Death Differ*, 14, 851-860 (2006)

61. Pardo, O. E., P. Wellbrock, U. Khanzada, M. Aubert, I. Arozarena, S. Davidson, F. Bowen, P. Parker, V. Filonenko, I. Gout, N. Sebire, R. Marais, J. Downward & M. Seckl: FGF-2 protects small cell lung cancer cells from apoptosis through a complex involving PKC, B-Raf and S6K2. *EMBO J*, 25, 3078-3088 (2006)

62. Steinberg, R., O. A. Harari, E. A. Lidington, J. J. Boyle, M. Nohadani, A. M. Samarel, M. Ohba, D. O. Haskard & J. C. Mason: A protein kinase C-epsilon-anti-apoptotic kinase signaling complex protects human vascular endothelial cells against apoptosis through induction of Bcl-2. *J Biol Chem*, 282, 32288-32297 (2007)

63. Ding, L., H. Wang, W. Lang & L. Xiao: Protein Kinase C-epsilon Promotes Survival of Lung cancer cells by suppressing apoptosis through dysregulation of the mitochondrial caspase pathway. *J Biol Chem*, 277, 35305-35313 (2002)

64. Mayne, G. C. & A. W. Murray: Evidence that protein kinase C-epsilon mediates phorbol ester inhibition of calphostin C- and tumor necrosis factor-alpha-induced

apoptosis in U937 histiocytic lymphoma cells. *J Biol Chem*, 273, 24115-24121 (1998)

65. Okhrimenko, H., W. Lu, C. Xiang, N. Hamburger, G. Kazimirsky & C. Brodie: Protein kinase C-epsilon regulates the apoptosis and survival of glioma cells. *Cancer Res*, 65, 7301-7309 (2005)

66. Shinohara, H., N. Kayagaki, H. Yagita, N. Oyaizu, M. Ohba, T. Kuroki & Y. Brodie: A protective role of PKC-epsilon against TNF-related apoptosis-inducing ligand (TRAIL)-induced apoptosis in glioma cells. *Biochem Biophys Res Commun*, 284, 1162-1167 (2001)

67. Koriyama, H., Z. Kouchi, T. Umeda, T. C. Saido, T. Momoi, S. Ishiura & K. Suzuki: Proteolytic activation of protein kinase C delta and epsilon by Caspase-3 in U937 cells during chemotherapeutic agent-induced apoptosis. *Cell Signal*, 11, 831-838 (1999)

68. Basu, A., D. Lu, B. Sun, A. N. Moor, G. R. Akkaraju & J. Huang: Proteolytic activation of Protein Kinase C-epsilon by caspase-mediated processing and transduction of antiapoptotic signals. *J Biol Chem*, 277, 41850-41856 (2002)

69. Castrillo, A., D. J. Pennington, F. Otto, P. J. Parker, M. J. Owen & L. Bosca: Protein Kinase C epsilon is required for macrophage activation and defense against bacterial infection. *J Exp Med*, 194, 1231-1242 (2001)

70. Akimoto, K., R. Takahashi, S. Moriya, N. Nishioka, J. Takayanagi, K. Kimura, Y. Fukui, S. Osada, K. Mizuno, S. Hirai, A. Kazlauskas & S. Ohno: EGF of PDGF receptors activate atypical PKC-lambda through phosphatidylinositol 3-kinase. *EMBO J*, 15, 788-798 (1996)

71. Diaz-Meco, M. T., E. Berra, M. M. Municio, L. Sanz, J. Lozano, I. Dominguez, V. Diaz-Golpe, M. T. Lain de Lera, J. Alcamí & C. V. Paya: A dominant negative protein kinase C-zeta subspecies blocks NF-kappa B activation. *Mol Cell Biol*, 13, 4770-4775 (1993)

72. Berra, E., M. T. Diaz-Meco, J. Lozano, S. Frutos, M. M. Municio, P. Sanchez, L. Sanz & J. Moscat: Evidence for a role of MEK and MAPK during signal transduction by protein kinase C zeta. *EMBO J*, 14, 6157-6163 (1995)

73. Filomenko, R., F. Poirson-Bichat, C. Billerey, J.-P. Belon, C. Garrido, E. Solary & A. Bettaieb: Atypical protein kinase C zeta as a target for chemosensitization of tumor cells. *Cancer Res*, 62, 1815-1821 (2002)

74. Leroy, I., A. de Thonel, G. Laurent & A. Quillet-Mary: Protein kinase C zeta associates with death inducing signaling complex and regulates Fas ligand-induced apoptosis. *Cell Signal*, 17, 1149-1157 (2005)

75. Plo, I., H. Hernandez, G. Kohlhagen, D. Lautier, Y. Pommier & G. Laurent: Overexpression of the atypical protein kinase C zeta reduces topoisomerase II catalytic activity, cleavable complexes formation, and drug-induced

- cytotoxicity in monocytic U937 leukemia cells. *J Biol Chem*, 277, 31407-31415 (2002)
76. Jamieson, L., L. Carpenter, T. J. Biden & A. P. Fields: Protein Kinase C- ι activity is necessary for Bcr-Abl-mediated resistance to drug-induced apoptosis. *J Biol Chem*, 274, 3927-3930 (1999)
77. Murray, N. R. & A. P. Fields: Atypical protein kinase C- ι protects human leukemia cells against drug-induced apoptosis. *J Biol Chem*, 272, 27521-4 (1997)
78. Garin, G., J.-i. Abe, A. Mohan, W. Lu, C. Yan, A. C. Newby, A. Rhaman & B. C. Berk: Flow antagonizes TNF- α signaling in endothelial cells by inhibiting caspase-dependent PKC- ζ processing. *Circ Res*, 101, 97-105 (2007)
79. Smith, L., L. Chen, M. E. Reyland, T. A. DeVries, R. V. Talanian, S. Omura & J. B. Smith: Activation of atypical protein kinase C- ζ by caspase processing and degradation by the ubiquitin-proteasome system. *J Biol Chem*, 275, 40620-40627 (2000)
80. Wang, G., J. Silva, K. Krishnamurthy, E. Tran, B. G. Condie & E. Bieberich: Direct binding to ceramide activates protein kinase C- ζ before the formation of a pro-apoptotic complex with PAR-4 in differentiating stem cells. *J Biol Chem*, 280, 26415-26424 (2005)
81. Berra, E., M. Municio, L. Sanz, S. Frutos, M. Diaz-Meco & J. Moscat: Positioning atypical protein kinase C isoforms in the UV-induced apoptotic signaling cascade. *Mol Cell Biol*, 17, 4346-4354 (1997)
82. LaVallie, E. R., P. S. Chockalingam, L. A. Collins-Racie, B. A. Freeman, C. C. Keohan, M. Leitges, A. J. Dorner, E. A. Morris, M. K. Majumdar & M. Arai: Protein Kinase C- ζ is up-regulated in osteoarthritic cartilage and is required for activation of NF- κ B by tumor necrosis factor and interleukin-1 in articular chondrocytes. *J Biol Chem*, 281, 24124-24137 (2006)
83. Diaz, J., T. Oltersdorf, W. Horne, M. McConnell, G. Wilson, S. Weeks, T. Garcia & L. C. Fritz: A common binding site mediates heterodimerization and homodimerization of Bcl-2 family members. *J Biol Chem*, 272, 11350-11355 (1997)
84. Diaz-Meco, M. T., M.-J. Lallena, A. Monjas, S. Frutos & J. Moscat: Inactivation of the inhibitory κ B protein kinase/nuclear factor κ B pathway by Par-4 expression potentiates tumor necrosis factor α -induced apoptosis. *J Biol Chem*, 274, 19606-19612 (1999)
85. Leitges, M., L. Sanz, P. Martin, A. Duran, U. Braun, J. F. Garcia, F. Camacho, M. T. Diaz-Meco, P. D. Rennert & J. Moscat: Targeted disruption of the ζ PKC gene results in the impairment of the NF- κ B pathway. *Mol Cell*, 8, 771-780 (2001)
86. Diaz-Meco, M., M. Municio, S. Frutos, P. Sanchez, J. Lazano, L. Sanz & J. Moscat: The product of *par-4*, a gene induced during apoptosis, interacts selectively with the atypical isoforms of protein kinase C. *Cell*, 86, 777-786 (1996)
87. Kim, S.-J., H.-G. Kim, C.-D. Oh, S.-G. Hwang, W.-K. Song, Y.-J. Yoo, S.-S. Kang & J.-S. Chun: p38 kinase-dependent and -independent inhibition of protein kinase C- ζ and - α regulates nitric oxide-induced apoptosis and dedifferentiation of articular chondrocytes. *J Biol Chem*, 277, 30375-30381 (2002)
88. Kim, J.-S., Z.-Y. Park, Y.-J. Yoo, S.-S. Yu & J.-S. Chun: p38 kinase mediates nitric oxide-induced apoptosis of chondrocytes through the inhibition of protein kinase C- ζ by blocking autophosphorylation. *J Biol Chem*, 280, 201-212 (2005)
89. Xin, M., F. Gao, W. S. May, T. Flagg & X. Deng: Protein Kinase C ζ abrogates the proapoptotic function of bax through phosphorylation. *J Biol Chem*, 282, 21268-21277 (2007)
90. Mao, M., X. Fang, Y. Lu, R. Lapushin, R. Bast & G. Mills: Inhibition of growth-factor-induced phosphorylation and activation of protein kinase B/Akt by atypical protein kinase C in breast cancer cells. *Biochem J*, 352, 475-482 (2000)
91. Doornbos, R. P., M. Theelen, P. C. J. van der Hoeven, W. J. van Blitterswijk, A. J. Verkleij & P. M. P. van Bergen en Henegouwen: Protein kinase C- ζ is a negative regulator of protein kinase B activity. *J Biol Chem*, 274, 8589-8596 (1999)
92. Powell, D. J., E. Hajdich, G. Kular & H. S. Hundal: Ceramide disables 3-phosphoinositide binding to the pleckstrin homology domain of protein kinase B (PKB)/Akt by a PKC- ζ -dependent mechanism. *Mol Cell Biol*, 23, 7794-7808 (2003)
93. Datta, R., H. Kojima, K. Yoshida & D. Kufe: Caspase-3-mediated cleavage of protein kinase C- θ in induction of apoptosis. *J Biol Chem*, 272, 20317-20 (1997)
94. Schultz, A., J.-I. Jönsson & C. Larsson: The regulatory domain of protein kinase C localises to the Golgi complex and induces apoptosis in neuroblastoma and Jurkat cells. *Cell Death Diff*, 10, 662-675 (2003)
95. Sun, X., F. Wu, R. Datta, S. Kharbanda & D. Kufe: Interaction between Protein Kinase C- δ and the c-Abl tyrosine kinase in the cellular response to oxidative stress. *J Biol Chem*, 275, 7470-7473 (2000)
96. Jackson, D., Y. Zheng, D. Lyo, Y. Shen, K. Nakayama, K. Nakayama, M. Humphries, M. Reyland & F. DA: Suppression of cell migration by protein kinase C- δ . *Oncogene*, 24, 3067-3072 (2005)

97. Santiago-Walker, A. E., A. J. Fikaris, G. D. Kao, E. J. Brown, M. G. Kazanietz & J. L. Meinkoth: Protein kinase C-delta stimulates apoptosis by initiating G1 phase cell cycle progression and S phase arrest. *J Biol Chem*, 280, 32107-32114 (2005)
98. Jackson, D. N. & D. A. Foster: The enigmatic protein kinase C-delta: complex roles in cell proliferation and survival. *FASEB J*, 18, 627-636 (2004)
99. Datta, K., R. Nambudripad, S. Pal, M. Zhou, H. T. Cohen & D. Mukhopadhyay: Inhibition of insulin-like growth factor-I-mediated cell signaling by the von Hippel-Lindau gene product in renal cancer. *J Biol Chem*, 275, 20700-20706 (2000)
100. Li, W., Y.-X. Jiang, J. Zhang, L. Soon, L. Flechner, V. Kapoor, J. H. Pierce & L.-H. Wang: Protein kinase C-delta is an important signaling molecule in insulin-like growth factor I receptor-mediated cell transformation. *Mol Cell Biol*, 18, 5888-5898 (1998)
101. Matassa, A., L. Carpenter, T. Biden, M. Humphries & M. Reyland: PKC-delta is required for mitochondrial dependent apoptosis in salivary epithelial cells. *J Biol Chem*, 276, 29719-29728 (2001)
102. Majumder, P., N. Mishra, X. Sun, A. Bharti, S. Kharbanda, S. Saxena & D. Kufe: Targeting of protein kinase C-delta to mitochondria in the oxidative stress response. *Cell Death Differ*, 12, 465-470 (2001)
103. Khwaja, A. & L. Tatton: Caspase-mediated proteolysis and activation of protein kinase C-delta plays a central role in neutrophil apoptosis. *Blood*, 94, 291-301 (1999)
104. Reyland, M., S. Anderson, A. Matassa, K. Barzen & D. Quissell: Protein kinase C-delta is essential for etoposide-induced apoptosis in salivary acinar cells. *J Biol Chem*, 274, 11915-11923 (1999)
105. Schechtman, D. & D. Mochly-Rosen: Isozyme-specific inhibitors and activators of protein kinase C. *Methods Enzymol*, 345, 470-489 (2002)
106. Reyland, M. E.: PKC and Apoptosis. In: *Apoptosis, Cell Signaling, and Human Disease: Molecular Mechanisms*. Ed: R. Srivastava. Humana Press Inc, Totowa, N.J. (2007)
107. Humphries, M. J., K. H. Limesand, J. C. Schneider, K. I. Nakayama, S. M. Anderson & M. E. Reyland: Suppression of apoptosis in the protein kinase C-delta null mouse *in vivo*. *J Biol Chem*, 281, 9728-9737 (2006)
108. Leitges, M., M. Mayr, U. Braun, U. Mayr, C. Li, G. Pfister, N. Ghaffari-Tabrizi, G. Baier, Y. Hu & Q. Xu: Exacerbated vein graft arteriosclerosis in protein kinase C delta-null mice. *J Clin Invest*, 108, 1505-1512 (2001)
109. Bharti, A., S.-K. Kraeft, M. Gounder, P. Pandey, S. Jin, Z.-M. Yuan, S. P. Lees-Miller, R. Weichselbaum, D. Weaver, L. B. Chen, D. Kufe & S. Kharbanda: Inactivation of DNA-dependent protein kinase by protein kinase C-delta: Implications for apoptosis. *Mol Cell Biol*, 18, 6719-6728 (1998)
110. Yoshida, K., H. Wang, Y. Miki & D. Kufe: Protein kinase C delta is responsible for constitutive and DNA damage induced phosphorylation of Rad9. *EMBO J*, 22, 1431-1441 (2003)
111. Voss, O. H., S. Kim, M. D. Wewers & A. I. Doseff: Regulation of monocyte apoptosis by the protein kinase C delta-dependent phosphorylation of caspase-3. *J Biol Chem*, 280, 17371-17379 (2005)
112. Sitailo, L. A., S. S. Tibudan & M. F. Denning: The protein kinase C-delta catalytic fragment targets Mcl-1 for degradation to trigger apoptosis. *J Biol Chem*, 281, 29703-29710 (2006)
113. Murriel, C. L., E. Churchill, K. Inagaki, L. I. Szweda & D. Mochly-Rosen: Protein kinase Cdelta activation induces apoptosis in response to cardiac ischemia and reperfusion damage: A mechanism involving BAD and the mitochondria. *J Biol Chem*, 279, 47985-47991 (2004)
114. Niwa, K., O. Inanami, T. Yamamori, T. Ohta, T. Hamasu, T. Karino & M. Kuwabara: Roles of protein kinase C-delta in the accumulation of p53 and the induction of apoptosis in H2O2-treated bovine endothelial cells. *Free Radic Res*, 36, 1147-1153 (2002)
115. DeVries, T. A., R. L. Kalkofen, A. A. Matassa & M. E. Reyland: Protein kinase C-delta regulates apoptosis via activation of STAT1. *J Biol Chem*, 279, 45603-45612 (2004)
116. Ren, J., R. Datta, H. Shioya, Y. Li, E. Oki, V. Biedermann, A. Bharti & D. Kufe: p73 beta is regulated by the protein kinase C delta catalytic fragment generated in the apoptotic response to DNA damage. *J Biol Chem*, 277, 33758-33765 (2002)
117. Yoshida, K., Y. Miki & D. Kufe: Activation of SAPK/JNK signaling by Protein Kinase C-delta in response to DNA damage. *J Biol Chem*, 277, 48372-48378 (2002)
118. Lee, Y.-J., D.-H. Lee, C.-K. Cho, S. Bae, G.-J. Jhon, S.-J. Lee, J.-W. Soh & Y.-S. Lee: HSP25 Inhibits Protein kinase Cdelta-mediated cell death through direct interaction. *J Biol Chem*, 280, 18108-18119 (2005)
119. Humphries, M., A. Ohm, J. Schaack, T. Adwan & M. Reyland: Tyrosine phosphorylation regulates nuclear localization of PKCdelta. *Oncogene*, In press, (2008)
120. DeVries-Seimon, T. A., A. M. Ohm, M. J. Humphries & M. E. Reyland: Induction of apoptosis is driven by nuclear retention of PKCdelta. *J Biol Chem*, 282, 22307-14 (2007)

121. Blass, M., I. Kronfeld, G. Kazimirsky, P. Blumberg & C. Brodie: Tyrosine phosphorylation of protein kinase C-delta is essential for its apoptotic effect in response to etoposide. *Mol Cell Biol*, 22, 182-195 (2002)
122. Konishi, H., M. Tanaka, Y. Takemura, H. Hiderori, Y. Ono, U. Kikkawa & Y. Nishizuka: Activation of protein kinase C by tyrosine phosphorylation in response to H2O2. *Proc Natl Acad Sci USA*, 94, 11233-11237 (1997)
123. Kharbanda, S., P. Pandey, S. Jin, S. Inoue, A. Bharti, Z.-M. Yuan, R. Weichselbaum, D. Weaver & D. Kufe: Functional interaction between DNA-PK and c-Abl in response to DNA damage. *Nature*, 386, 732-735 (1997)
124. Zang, Q., Z. Lu, M. Curto, N. Barile, D. Shalloway & D. A. Foster: Association between v-Src and protein kinase C-delta in v-Src-transformed fibroblasts. *J Biol Chem*, 272, 13275-13280 (1997)
125. DeVries, T. A., M. C. Neville & M. E. Reyland: Nuclear import of PKC-delta is required for apoptosis: Identification of a novel nuclear import sequence. *EMBO J*, 21, 6050-6060 (2002)
126. Yuan, Z.-M., T. Utsugisawa, T. Ishiko, S. Nakada, Y. Huang, S. Kharbanda, R. Weichselbaum & D. Kufe: Activation of protein kinase C-delta by the c-Abl tyrosine kinase in response to ionizing radiation. *Oncogene*, 16, 1643-1648 (1998)
127. Sitailo, L. A., S. S. Tibudan & M. F. Denning: Bax activation and induction of apoptosis in human keratinocytes by the protein kinase C delta catalytic domain. *J Invest Dermatol*, 123, 434-443 (2004)
128. Castanga, M., Y. Takai, K. Kaibuchi, K. Sano, U. Kikkawa & Y. Nishizuka: Direct activation of calcium-activated, phospholipid dependent protein kinase by tumor-promoting phorbol esters. *J Biol Chem*, 257, 7847-7851 (1982)
129. Kikkawa, U., Y. Takai, Y. Tanaka, R. Miyake & Y. Nishizuka: Protein kinase C as a possible receptor protein of tumor-promoting phorbol esters. *J Biol Chem*, 258, 11442-11445 (1983)
130. Lu, Z., A. Hornia, Y.-W. Jiang, Q. Zang, S. Ohno & D. Foster: Tumor promotion by depleting cells of protein kinase C-delta. *Mol Cell Biol*, 17, 3418-3428 (1997)
131. Reno, E. M., J. M. Haughian, I. K. Dimitrova, T. A. Jackson, K. R. Shroyer & A. P. Bradford: Analysis of protein kinase C delta expression in endometrial tumors. *Human Pathology*, 39, 21-29 (2008)
132. D'Costa, A., J. Robinson, T. Maududi, V. Chaturvedi, B. Nickoloff & D. M.F.: The proapoptotic tumor suppressor protein kinase C-delta is lost in human squamous cell carcinomas. *Oncogene*, 25, 375-386 (2006)
133. Tan, M., P. Li, M. Sun, G. Yin & D. Yu: Upregulation and activation of PKC by ErbB2 through Src promotes breast cancer cell invasion that can be blocked by combined treatment with PKC and Src inhibitors. *Oncogene*, 25, 3286-3295 (2006)
134. Michie, A. M. & R. Nakagawa: The link between PKC-alpha regulation and cellular transformation. *Immunol Lett*, 96, 155-162 (2005)
135. Lahna, M., G. Köhlerd, K. Sundella, C. Sub, S. Lib, B. Paterson & T. Bumolc: Protein kinase C -alpha expression in breast and ovarian cancer. *Oncology*, 67, 1-10 (2004)
136. Jiang, X.-H., S.-P. Tu, J.-T. Cui, M. C. M. Lin, H. H. X. Xia, W. M. Wong, A. O.-O. Chan, M. F. Yuen, S.-H. Jiang, S.-K. Lam, H.-F. Kung, J. W. Soh, I. B. Weinstein & B. C.-Y. Wong: Antisense targeting protein kinase C-alpha and -beta1 inhibits gastric carcinogenesis. *Cancer Res*, 64, 5787-5794 (2004)
137. Shen, L., N. Dean & R. Glazer: Induction of p53-dependent, insulin-like growth factor-binding protein-3-mediated apoptosis in glioblastoma multiforme cells by a protein kinase C-alpha antisense oligonucleotide. *Mol Pharmacol*, 55, 396-402 (1999)
138. Lahn, M., K. Sundell & S. Moore: Targeting protein kinase C-alpha in cancer with the phosphorothioate antisense oligonucleotide aprinocarsen. *Ann NY Acad Sci*, 1002, 263-270 (2003)
139. Grossman, S., J. Alavi, J. Supko, K. Carson, R. Priet, F. Dorr, J. Grundy & J. Holmlund: Efficacy and toxicity of the antisense oligonucleotide aprinocarsen directed against protein kinase C- alpha delivered as a 21-day continuous intravenous infusion in patients with recurrent high-grade astrocytomas. *Neuro-Oncology*, 1, 32-40 (2005)
140. Vansteenkiste, J., J.-L. Canon, H. Riska, R. Pirker, P. Peterson, W. John, P. Mali & M. Lahn: Randomized phase II evaluation of aprinocarsen in combination with gemcitabine and cisplatin for patients with advanced/metastatic non-small cell lung cancer. *Invest New Drugs*, 23, 263-269 (2005)
141. Pan, Q., L. W. Bao, C. G. Kleer, M. S. Sabel, K. A. Griffith, T. N. Teknos & S. D. Merajver: Protein kinase C-epsilon is a predictive biomarker of aggressive breast cancer and a validated target for RNA interference anticancer therapy. *Cancer Res*, 65, 8366-8371 (2005)
142. Knauf, J. A., R. Elisei, D. Mochly-Rosen, T. Liron, X.-N. Chen, R. Gonsky, J. R. Korenberg & J. A. Fagin: Involvement of protein kinase Cε in thyroid cell death. A truncated chimeric PKC-epsilon cloned from a thyroid cancer cell line protects thyroid cells from apoptosis. *J Biol Chem*, 274, 23414-23425 (1999)
143. Sharif, T. & M. Sharif: Overexpression of protein kinase C-epsilon in astroglial brain tumor derived cell lines

and primary tumor samples. *Int J Oncol*, 15, 237-243 (1999)

144. Cornford, P., J. Evans, A. Dodson, K. Parsons, A. Woolfenden, J. Neoptolemos & C. S. Foster: Protein kinase C isoenzyme patterns characteristically modulated in early prostate cancer. *Am J Pathol*, 154, 137-144 (1999)

145. Wu, D., T. L. Foreman, C. W. Gregory, M. A. McJilton, G. G. Wescott, O. H. Ford, R. F. Alvey, J. L. Mohler & D. M. Terrian: Protein kinase C-epsilon has the potential to advance the recurrence of human prostate cancer. *Cancer Res*, 62, 2423-2429 (2002)

146. McJilton, M.A., C. Van. Sikes, G.G. Wescott, D. Wu, T.L, Foreman, C.W. Gregory, D.A. Weidner, O. H. Ford, A. M. Lasater, J.L. Mohler & D.M. Terrian: Protein kinase C-epsilon interacts with Bax and promotes survival of human prostate cancer cells. *Oncogene*, 22, 7958-7968 (2003)

147. Tachado, S., M. Mayhew, G. Wescott, T. Foreman, C. Goodwin, M. McJilton & D. Terrian: Regulation of tumor invasion and metastasis in protein kinase C-epsilon-transformed NIH3T3 fibroblasts. *J Cell Biochem*, 85, 785-797 (2002)

148. Griner, E. M. & M. G. Kazanietz: Protein kinase C and other diacylglycerol effectors in cancer. *Nat Rev Cancer*, 7, 281-294 (2007)

Abbreviations: protein kinase C: PKC, diacylglycerol: DAG, Receptors for Activated C Kinase: RACK's, Signal Transducer and Activator of Transcription 3: STAT 3, Prostate Apoptosis Response-4: PAR-4, Prostate Apoptosis Response-6: PAR-6, kinase inactive PKC-delta: PKC-delta KD, Signal Transducer and Activator of Transcription 1:STAT1, PKC-delta catalytic fragment:delta-CF

Key Words: Protein kinase C, Phosphorylation, Apoptosis, Proliferation, Survival, Cancer, Review

Send correspondence to: Mary E. Reyland, Department of Craniofacial Biology, University of Colorado HSC, Mail Stop 8120, P.O. Box 6511, Aurora, CO 80045, Tel: 303-724-4572, Fax:303 724-4580, E-mail: Mary.Reyland@UCHSC.edu