

Endocytosis and membrane receptor internalization: implication of F-BAR protein Carom

Yanjie Xu^{1,2}, Jixiang Xia², Suxuan Liu^{2,6}, Sam Stein², Cueto Ramon², Hang Xi², Luqiao Wang^{1,2}, Xinyu Xiong², Lixiao Zhang², Dingwen He⁷, William Yang², Xianxian Zhao⁶, Xiaoshu Cheng¹, Xiaofeng Yang^{2,3,4,5}, Hong Wang^{2,3,4,5}

¹Center Department of Cardiology, Second Affiliated Hospital of Nanchang University, Nan Chang, Jiang Xi, 330006, China, ²Center for Metabolic Disease Research, Temple University School of Medicine, Philadelphia, PA, 19140, ³Cardiovascular Research, Temple University School of Medicine, Philadelphia, PA, 19140, ⁴Thrombosis Research, Temple University School of Medicine, Philadelphia, PA, 19140, ⁵Department of Pharmacology, Temple University School of Medicine, Philadelphia, PA, 19140, ⁶Department of Cardiology, Changhai Hospital, Second Military Medical University, Shanghai, 200433, China, ⁷Department of Orthopedics, Second Affiliated Hospital of Nanchang University, Nan Chang, Jiang Xi, 330006, China

TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Materials and Methods
 - 3.1. Identification of endocytic membrane receptor (Kegg database, Genecard database)
 - 3.2. Identification of regulating protein in CDE, F-BAR protein potential in membrane receptor endocytosis and Carom potential endocytic partners (Pubmed, Genecard database)
 - 3.3. Cell expression profile of endocytic membrane receptor, Carom and Carom endocytic partner (Genevestigator database)
 - 3.4. Predicted interaction of endocytic membrane receptor with Carom and its endocytic partners. (String database, NCBI database)
 - 3.5. Subcellular localization analysis of Carom endocytic partner and membrane receptor, and co-localization with Carom (Compartments database, cNLS Mapper)
4. Results
 - 4.1. Identification of 34 endocytic membrane receptors.
 - 4.2. Key steps in vesicle formation in CDE and its regulating proteins
 - 4.3. F-BAR protein participates in membrane receptor endocytosis
 - 4.4. Identification of 9 Carom endocytic partners
 - 4.5. Endocytic membrane receptor, Carom and its endocytic partner are differently expressed in human primary cells
 - 4.6. Carom may directly or indirectly bind to endocytic membrane receptor through partner protein
 - 4.7. Carom co-localization with endocytic membrane receptors and partners in different cellular micro-compartment
 - 4.8. Carom-receptor complex intracellular trafficking and function
5. Discussion
6. Conclusion
7. Acknowledgements
8. References

1. ABSTRACT

Endocytosis is a cellular process mostly responsible for membrane receptor internalization. Cell membrane receptors bind to their ligands and form a complex which can be internalized. We

previously proposed that F-BAR protein initiates membrane curvature and mediates endocytosis via its binding partners. However, F-BAR protein partners involved in membrane receptor endocytosis and the




Figure 1. Overview of endocytosis. Endocytosis is a cellular process by which molecules or substances are transported into the cell via cell membrane engulfment. A. Classification of endocytosis. Endocytosis is generally classified as phagocytosis and pinocytosis. Pinocytosis can be further divided into 4 subtypes; macropinocytosis, clathrin-dependent, caveolae-dependent, and clathrin/caveolae independent endocytosis based on clathrin or caveolae involvement. Most of the receptor-mediated endocytosis (REM) is processed via clathrin-dependent mechanism. B. Features of endocytosis. Features of endocytosis are summarized for the size of the internalized vesicle, membrane domain localization and cargo content. C. Schematic diagram of endocytosis process. Pathogens and ligands induce endocytosis by binding to the cell membrane via receptor-dependent or -independent mechanism, and then form phagosome or endocytic vesicle which may be coated with clathrin/caveolin or regulated by flotillin, GRAF1, Arf6 and RhoA. Membrane-bounded F-BAR protein is linked to actin-associated proteins, can cause cell membrane curvature and facilitate clathrin-mediated or caveolae-dependent endocytosis. During phagocytosis, cells bring in solid particles into phagosomes and then fuse with lysosomes (marked by LAMP1). During pinocytosis, internalized vesicles are transported to early endosome (marked by Rab5, EEA1). The early endosome can send the cargoes to three locations: 1) late endosome (marked by Rab7) then lysosome for degradation, 2) recycling endosome (marked by Rab4,11) for signal transduction or recycling to plasma membrane, and 3) nucleus for transcription factor regulation or chromatin remodeling machinery. Abbreviation: RTK, receptor tyrosine kinase; GPCR, G protein-coupled receptor; TFR, transferrin receptor; LDLR, low-density lipoprotein receptor; GPI, glycosylphosphatidylinositol; TGF-βR, transforming growth factor-beta receptor; IGF-1R, insulin-like growth factor I receptor; IL-2RB, interleukin 2 receptor beta; Rab, Ras associated protein; EEA1, early endosome antigen 1; LAMP1, lysosomal associated membrane protein 1; GRAF1, Rho GTPase Activating Protein 26; Arf6, ADP-Ribosylation Factor 6; RhoA, Ras Homolog Family Member A.

regulatory mechanism remain unknown. In this study, we established database mining strategies to explore mechanisms underlying receptor-related endocytosis. We identified 34 endocytic membrane receptors and 10 regulating proteins in clathrin-dependent endocytosis (CDE), a major process of membrane receptor internalization. We found that F-BAR protein FCHSD2 (Carom) may facilitate endocytosis via 9 endocytic partners. Carom is highly expressed, along with highly expressed endocytic membrane receptors and partners, in endothelial cells and macrophages. We established 3 models of Carom-receptor complexes and their intracellular trafficking based on protein interaction and subcellular localization. We conclude that Carom may mediate receptor endocytosis and transport endocytic receptors to the cytoplasm for receptor signaling and lysosome/proteasome degradation, or to the nucleus for RNA processing, gene transcription and DNA repair.

2. INTRODUCTION

Endocytosis is a cellular process by which molecules or substances are transported into the cell via cell membrane engulfment. Endocytosis is generally classified as phagocytosis and pinocytosis, which are distinguished by the size of the endocytic vesicles formed (Figure 1) (1). Phagocytosis implies the ingestion of large and solid particles (diameter 0.5-10µm) such as pathogens. Pinocytosis refers to internalization of various liquids via small endocytic vesicles and can be divided into four subtypes: macropinocytosis, clathrin-dependent, caveolae-dependent, and clathrin/caveolae-independent endocytosis based on clathrin or caveolae involvement (2). Pathogens or ligands induce endocytosis by binding to the cell membrane via receptor-dependent or -independent mechanisms, and then form phagosome or endocytic vesicle (Figure 1). Endocytic vesicle




Figure 2. Key steps in clathrin-dependent endocytosis and its regulating proteins. A. Key steps in vesicle formation in clathrin-dependent endocytosis (CDE). There are four steps during vesicle formation: (1) curvature initiation, (2) vesicle formation, (3) vesicle scission, and (4) un-coating vesicle. At first, F-BAR protein binds to the plasma membrane and initiates membrane curvature. F-BAR protein can recruit adaptor protein via its SH3 domain during vesicle formation. Clathrin is recruited directly from the cytosol to the site of adaptor-concentrated membrane to help the formation of coated vesicle. GTPase dynamin can then bind to the membrane and cause vesicle constriction, scission, and release. HSC70 binds to clathrin, disassociates clathrin, intersectin and dynamin from the vesicle and produces an un-coated endocytic vesicle containing the cargo molecules. B. Regulating proteins during vesicle formation in CDE. A group of proteins are involved in endocytic vesicle formation. F-BAR protein initiates membrane curvature and clathrin-coated endocytic vesicle formation. Adaptor proteins (Intersectin, AP2, Epsin, CALM) link various components of the clathrin machinery to the membrane and help the formation of adaptor-concentrated clathrin-coated vesicle. Dynamin triggers vesicle scission upon GTP hydrolysis. HSC70 triggers un-coating of endocytic vesicle. Abbreviation: AP2, adaptor protein 2; SNX9, sorting nexin 9; HSC70, ATPase heat shock cognate 70; CALM, clathrin assembly lymphoid myeloid leukaemia; (N)-WASP/WAVE, Wiskott-Aldrich Syndrome Like.

may be coated with clathrin, caveolin or regulated by flotillin, Rho GTPase activating protein 26 (GRAF1), ADP-Ribosylation factor 6 (Arf6) and Ras homology family membrane A (RhoA). During phagocytosis, solid particle containing-phagosomes fuse with lysosomes (marked by lysosomal associated membrane protein (LAMP1)) and undergo lysosomal degradation. In the process of pinocytosis, internalized vesicles are transported to the early endosome (marked by Ras associated protein (Rab5)), which delivers the cargoes to three locations: 1) late endosome (marked by Ras associated protein (Rab7)) then lysosome for degradation, 2) recycling endosome (marked by Rab11) for signal transduction or receptor recycling to cell membrane, and 3) nucleus to regulate transcription factor and chromatin remodeling (1-3).

Membrane receptors are responsible for transducing external signals into the cell by receiving extracellular molecules. It is suggested that some of the cell membrane receptors bind to their ligands and form a complex which can be internalized and translocated to the cytoplasm or nucleus for signaling or degradation mostly via clathrin-dependent endocytosis (CDE) mechanism (4). For example, receptor tyrosine kinases

(RTKs) and G-protein coupled receptors (GPCRs) can be internalized by directly interacting with adaptor protein AP2, epsin or intersectin and CDE-mediated mechanism (5, 6). However, molecular mechanisms regulating CDE signaling are not fully elucidated.

We previously proposed that F-BAR (Fes/CIP4 homology-BAR) protein initiates membrane curvature and mediates endocytosis via its binding partners (7, 8). Beyond F-BAR domain which initiates endocytosis, most of the proteins in this superfamily contain other domains, such as Src homology-3 (SH3) and Src homology-2 (SH2) domain, which can recruit adaptor proteins to form complexes. Such F-BAR protein complexes participate in multiple steps of endocytosis, ranging from the assembly of endocytic vesicles and their scissions, F-actin polymerization and nucleation, etc. As a novel member of F-BAR protein, FCHSD2 (Carom) displays such structures and contains a F-BAR domain and two SH3 domains. Although this protein's function hasn't been fully addressed, it is predicted as a critical molecule in the process of CDE (Figure 2). However, how Carom interacts with membrane receptor and facilitate receptor endocytosis is unknown.

In the past ten years, bioinformatics analysis has emerged as an important tool for functional interpretation of genomics and proteomics information (9-11). In this study, we established a group of database mining strategies and performed intensive literature searches to explore mechanisms underlying receptor-related endocytosis. We identified endocytic membrane receptors and potential regulating proteins for vesicle formation and investigated the relationship of F-BAR protein Carom with endocytic membrane receptors and endocytic partners. We established models of endocytosis and Carom-mediated membrane receptor internalization.

3. MATERIALS AND METHODS

3.1. Identification of endocytic membrane receptor (Kegg database, Genecard database)

Thirty-four cell membrane receptors were selected from 259 genes related to endocytosis identified from Kegg pathway database (hsa04144, <http://www.genome.jp/dbget-bin/www.bget?hsa04144>) (Table 1). The ligands and function of receptors were identified using Genecard database (<http://www.genecards.org>). Endocytosis type was defined by literature search.

3.2. Identification of regulating protein in CDE, F-BAR protein potential in membrane receptor endocytosis and Carom potential endocytic partners (Pubmed, Genecard database)

We searched through PubMed literature to summarize 10 important regulating proteins that take part in vesicle formation in CDE and F-BAR protein potential in membrane receptor endocytosis (Figure 2, Table 2). A total of 26 binding partners for Carom protein were identified from previous study, which was established via Affinity Capture-MS, Affinity Capture-RNA, Affinity Capture-Western, Reconstituted Complex and Two-hybrid technologies (12, 13). 9 of 26 Carom endocytic partners were selected based on literature review and their function (13-15) (Table 3). Functions and binding domains of Carom partners were defined based on information obtained from Genecard database. Binding receptors were identified from NCBI Gene database and their corresponding regulation on receptors was determined by PubMed search (16-18).

3.3. Cell expression profile of endocytic membrane receptor, Carom and Carom endocytic partner (Genevestigator database)

To investigate the expression profile of the endocytic membrane receptors, Carom and its endocytic partners in the cells, we used bioinformatics methods to gather extensive microarray information in the human primary cells. mRNA levels were obtained from microarray data available on the web site (<https://www.genevestigator.com/gv/>) and expressed as a heat

map (19) (Figure 3). The dark and light color shading represents the relatively high and low expression levels respectively in the different human primary cells. The three darker shadings represent higher levels of expression.

3.4. Predicted interaction of endocytic membrane receptor with Carom and its endocytic partners. (String database, NCBI database)

To generate an overview of whether Carom and its 9 endocytic partners may relate to membrane receptor functionality, we carried out an analysis of their protein-protein interactions by using String database (19, 20) (Figure 4). Solid lines indicate known interaction deposited in NCBI Gene database, which was established based on Affinity Capture-MS, Affinity Capture-RNA, Affinity Capture-Western, Reconstituted Complex and Two-hybrid experimental data. Dashed lines indicate computational-predicted interaction in String database based on analyzing genomic information ('genomic context'-methods) or from transferring associations/interactions between organisms ('interolog'-transfer).

3.5. Subcellular localization analysis of Carom endocytic partner and membrane receptor, and co-localization with Carom (Compartments database, cNLS Mapper)

Subcellular localization of Carom, its endocytic partner and membrane receptors were determined using Compartments database established by manually curated literature, high-throughput screens, automatic text mining, and sequence-based prediction methods (20) (Table 4). Numbers are indicated confidence levels. Protein nuclear import generally involves the nuclear localization signal (NLS), or interaction with carrier proteins (21). Nuclear localization signal was identified by analyzing protein sequence in cNLS Mapper. We predicted Carom-receptor complex intracellular trafficking from cell membrane into cytoplasm based on the analysis of co-localization of endocytic membrane receptor and partner in plasma membrane (PM), cytoplasm (CP) and cytoskeleton (CSK). We predicted Carom-receptor complex translocation into the nucleus (NC) based on the analysis of co-localization of endocytic membrane receptor and partner in PM, NC and NLS. We predicted the function of Carom-receptor complexes based on receptor signaling reported in the literature (Table 5)

4. RESULT

4.1. Identification of 34 endocytic membrane receptors

We identified 34 membrane receptors from 259 endocytosis-related genes selected from Kegg pathway database and classified them into three

Table 1. Classification of endocytic membrane receptor

Gene Symbol Full name)	Ligand	Function	Endocytosis type
G-protein coupled receptor (GPCR)			
1. ADRB1 (Adrenoceptor β 1)	Epinephrine, norepinephrine	Mediate catecholamines action	CDE
2. ADRB2 (Adrenoceptor β 2)	Epinephrine, norepinephrine	Mediate catecholamines action	CDE
3. ADRB3 (Adrenoceptor β 3)	Norepinephrine	Mediate catecholamines action	CDE
4. CCR5 (Chemokine (C-C motif) receptor 5)	CCl3, CCl4, CCl5, CCl8, CCl13, CCl16	Leuc trafficking, angiogenesis, apoptosis	CDE
5. CXCR1 (Chemokine (C-X-C motif) receptor 1)	CXCl6, CXCl8	Leuc trafficking, angiogenesis, apoptosis	CDE
6. CXCR2 Chemokine (C-X-C motif) receptor 2)	CXCl1, CXCl2, CXCl3, CXCl5, CXCl6, CXCl7, CXCl8	Leuc trafficking, angiogenesis, apoptosis	CDE
7. CXCR4 (Chemokine (C-X-C motif) receptor 4)	CXCl14	Leuc trafficking, angiogenesis, apoptosis	CDE
8. F2R (Coagulation factor II receptor)	Thrombin	PLT activation, vascular development	CDE
Receptor tyrosine kinase (RTK)			
9. CSF1R (Colony stimulating factor 1 receptor)	M-CSF, IL34	Macrophage regulator	CDE
10. EGFR (Epidermal growth factor receptor)	EGF	Proliferation, differentiation	CDE/CIE
11. ERBB2 (Erb-b2 receptor tyrosine kinase 2)	EGF	Proliferation, differentiation	CDE
12. ERBB3 (Erb-b2 receptor tyrosine kinase 3)	EGF	Proliferation, differentiation	CDE
13. ERBB4 (Erb-b2 receptor tyrosine kinase 4)	EGF	Proliferation, differentiation	CDE
14. FGFR1 (Fibroblast growth factor receptor 1)	FGF1, FGF2, FGF3, FGF6, FGF7	Proliferation, differentiation	CDE/CIE
15. FGFR2 (Fibroblast growth factor receptor 2)	FGF1, FGF4, FGF6, FGF7, FGF8	Proliferation, differentiation	CDE/CIE
16. FGFR3 (Fibroblast growth factor receptor 3)	FGF3, FGF4, FGF5, FGF6, FGF7	Proliferation, differentiation	CDE/CIE
17. FGFR4 (Fibroblast growth factor receptor 4)	FGF1, FGF3, FGF4, FGF5, FGF9	Proliferation, differentiation	CDE/CIE
18. FLT1 (Fms-related tyrosine kinase 1)	VEGFA, VEGFB, PGF	Angiogenesis	CDE/CIE
19. IGF1R (Insulin-like growth factor 1 receptor)	IGF1, IGF2	Proliferation, differentiation	CDE/CIE
20. IGF2R (Insulin-like growth factor 2 receptor)	IGF2, Transferrin	Proliferation, differentiation	CDE
21. KDR (Kinase insert domain receptor)	VEGFA, VEGFC	Proliferation, angiogenesis	CDE/CIE
22. MET (Tyrosine-protein kinase met)	HGF	Proliferation, angiogenesis	CDE
23. NTRK1 (Neurotrophic tyrosine kinase receptor type 1)	NGF	Differentiation	CDE
24. PDGFRA (Platelet-derived growth factor α receptor)	PDGFC	Proliferation, differentiation	CDE
25. TGFBR1 (Transforming growth factor β receptor I)	TGF- β	Proliferation, tumor transformation	CDE/CIE
26. TGFBR2 (Transforming growth factor β receptor I)	TGF- β	Proliferation, tumor transformation	CDE/CIE
Transmembrane receptor (TMR)			
27. FOLR1 (Folate receptor 1)	Folic acid	Transport folic acid	CDE
28. FOLR2 (Folate receptor 2)	Folic acid	Transport folic acid	CDE
29. FOLR3 (Folate receptor 3)	Folic acid	Transport folic acid	CDE
30. IL2RA (Interleukin 2 receptor α)	IL2	Regulate immune system	Indt
31. IL2RB (Interleukin 2 receptor β)	IL2, IL15	Regulate immune system	Indt
32. IL2RG (Interleukin 2 receptor γ)	IL2, IL-4, IL15	Regulate immune system	Indt
33. LDLR (Low density lipoprotein receptor)	LDL, ApoB100, ApoE, IDL	Transport lipid	CDE
34. TFRC (Transferrin receptor)	Transferrin, HFE	Transport iron	CDE

Classification of endocytic membrane receptor. We selected 34 cell surface receptors from 259 genes related to endocytosis identified from Kegg pathway database (hsa04144, <http://www.genome.jp/dbget-bin/www.bget?hsa04144>) and classified them into three groups: 1) G-protein coupled receptor, 2) Receptor tyrosine kinase, and 3) Transmembrane receptor. Ligands and function of receptors were identified using Genecard database (<http://www.genecards.org>). Endocytosis type was defined by literature search. Noted that most of the receptor-mediated endocytosis generally occurs via CDE. Abbreviation: Leuc, leukocyte; PLT, platelets; CDE, clathrin-dependent endocytosis; CIE, clathrin-independent endocytosis; Indt, clathrin/caveolae-independent endocytosis.

Table 2. F-BAR proteins are involved in membrane receptor endocytosis

F-BAR Proteins	Endocytosis	Roles in endocytosis	PMID#
1. CIP4 subfamily			
CIP4	√	Required for EGFR trafficking /degradation	19632321
FBP17	√	Required for EGFR internalization	19632321
Toca-1	√	Required for EGFR trafficking from endosomes	19632321
2. FCHOs subfamily			
FCHO1	√	Forming clathrin-coated vesicle	20448150
FCHO2	√	Required for LDLR endocytosis	22323290
3. srGAPs subfamily			
srGAP1	N/A		
srGAP2	N/A		
srGAP3	N/A		
4. PACSINs subfamily			
PACSIN1	√	Inhibiting endocytosis	11082044
PACSIN2	√	Required for EGFR translocated to endosomes	23129763
PACSIN3	√	Inhibiting transferrin/TFRC complex endocytosis	11082044
5. PSTPIPs subfamily			
PSTPIP1	√	Suppressing transferrin/TFRC complex endocytosis	18480402
PSTPIP2	N/A		
6. FCHSDs subfamily			
FCHSD1	√	Promoting F-actin polymerization and facilitate endocytosis	23437151
FCHSD2 (Carom)	√	Stimulating F-actin polymerization and facilitate endocytosis	23437151
7. FES/FER subfamily			
FES	N/A		
FER	N/A		
8. NOSTRIN subfamily	√	Assembling NOSTRIN-FGFR1-Rac1-Sos1 complex/regulate FGF signaling	22751148
9. GAS7 subfamily	N/A		

F-BAR proteins are involved in membrane receptor endocytosis. F-BAR proteins have 9 family members, each protein contains one F-BAR domain and other domains such as SH3, SH2, WW, and RhoGAP. Most of the F-BAR proteins are involved in endocytosis and play important roles in membrane receptor trafficking/signaling cited by PMID#. Abbreviation: F-BAR, Fes/CIP4 homology-BAR; CIP4, Cdc42-interacting protein 4; FBP17, Formin binding protein 17; Toca-1, TOCA homolog 1; FCHO1-2, FCH domain only 1-2; srGAP1-3, SLIT-ROBO Rho GTPase activating protein 1-3; PSTPIP1-2, proline-serine-threonine phosphatase-interacting protein 1-2; FCHSD1-2, FCH and double SH3 domains 1-2; NOSTRIN, nitric oxide synthase traffic inducer; GAS7, growth arrest specific 7; Others refer to table 1.

groups (Table 1). 1) 8 G-protein coupled receptors (GPCR): Adrenoceptor β 1-3 (ADRB1-3), Chemokine C-C motif receptor 5 (CCR5), Chemokine C-X-C motif 1,2,4 (CXCR1,2,4) and Coagulation factor II receptor (F2R), 2) 18 tyrosine kinase receptors (RTK): Colony stimulating factor 1 receptor (CSF1R), Epidermal growth factor receptor (EGFR), Erb-b2 receptor tyrosine kinase 2-4 (ERBB2-4), Fibroblast growth factor receptor 1-4 (FGFR1-4), Fms-related tyrosine kinase 1/Vascular endothelial growth factor receptor 1 (FLT1 or VEGFR1), Insulin-like growth factor 1-2 receptor (IGF1-2R), Kinase insert domain receptor/Vascular endothelial growth factor receptor 2 (KDR or VEGFR2), Tyrosine-protein kinase met (MET), Neurotrophic tyrosine kinase receptor type 1 (NTRK1),

Platelet-derived growth factor α receptor (PDGFRA), Transforming growth factor β receptor 1-2 (TGFB1-2), and 3) 8 transmembrane receptors (TRM): Folate receptor 1-3 (FOLR1-3), Interleukin 2 receptor α , β , γ (IL2RA, B, G), Low density lipoprotein receptor (LDLR) and Transferrin receptor (TFRC). It is known that endocytic membrane receptors can be internalized upon binding to their ligands and regulate various cellular functions including angiogenesis, proliferation, differentiation, and lipid/iron transport. The majority of these endocytic membrane receptors are internalized via CDE mechanism (22). Some of the receptors, such as EGFR, FGFR1-4, FLT1 (VEGFR1), IGF1R and TGFB1-2 can be internalized through both CDE and clathrin-independent endocytosis (CIE) (23). IL2R

Table 3. Identification of F-BAR protein Carom endocytic partner involved in endocytosis

Carom endocytic partner Symbol Full name	Function	Domains on partner	Binding Receptor	Receptor regulation
1. DAPP1 (B Lymphocyte Adapter Protein Bam32)	Regulates B-cell antigen receptor signaling	SH2, PH	ERBB3	N/A
2. GRASP (GRP1-Associated Scaffold Protein)	Regulate intracellular trafficking	PDZ	N/A	N/A
3. ITSN1 (Intersectin 1)	Regulate endocytic vesicle Formation	EF, DH, PH, C2	N/A	EGFR ubiquitination
4. ITSN2 (Intersectin 2)	Regulate endocytic vesicle Formation	EF, DH, PH, C2	EGFR	N/A
5. UBC (Ubiquitin C)	Regulate protein ubiquitination	Ubiquitin	EGFR	EGFR ubiquitination
6. UBD (Ubiquitin D)	Regulate protein ubiquitination	Ubiquitin	N/A	EGFR ubiquitination
7. VCP (Valosin Containing Protein)	Regulate vesicle trafficking	AAA ATPase	N/A	EGFR degradation
8. WAS (Wiskott-Aldrich Syndrome)	Regulate actin polymerization	WH1/2, PH, CRIB	EGFR	N/A
9. WASL (Wiskott-Aldrich Syndrome Like)	Regulate actin polymerization	WH1/2, PH, CRIB	EGFR	EGFR endocytosis

Identification of F-BAR protein Carom endocytic partner involved in endocytosis. Nine Carom endocytic partners were selected based on literature review from 26 Carom partners we identified previously (Front Biosci (Landmark Ed), 2016; 21:856-72), which were established via Affinity Capture-MS, Affinity Capture-RNA, Affinity Capture-Western, Reconstituted Complex and Two-hybrid technologies. Functions and binding domains of Carom partners were defined based on information obtained from Genecard database. Binding receptors were identified from NCBI Gene database and their regulation on receptor were determined by PubMed search. All the Carom partners contain different domains such as SH2, PH, PDZ, EF and Ubiquitin, which can recruit proteins to organize signaling complexes at cellular membranes. Noted that DAPP1 can bind to ERBB3, ITSN2/UBC/WAS/WASL can bind to EGFR, and that ITSN1/UBC/UBD, VCP and WASL can regulate EGFR ubiquitination, degradation and endocytosis process. Abbreviation: CRIB, CDC42-Rac interactive binding; EF, EF hand; DH, DBL homology; AAA ATPase, ATPases associated with diverse cellular activities; PH, Pleckstrin homology; PDZ, PSD-95/Dlg-AZO-1; WH1/2, WASP-Homology 1/2; Others refer to table 1.




Figure 3. Endocytic membrane receptor, Carom and Carom endocytic partner expression profile in human primary cells. A. Heat map of membrane receptor, Carom and Carom partner expression in human primary cells. mRNA levels are obtained from microarray data available in the web site (<https://www.geneinvestigator.com/gv/>) and expressed as a heat map. Gradient bars indicate percent of expression potential. The dark and light color shadings represent relatively high and low expression levels respectively. Dashed frames indicate body systems and cells with relatively high gene expression. B. Relationship of Carom, endocytic membrane receptor and partner expression in human circulatory and immune system cells. Noted that in aortic ECs, membrane receptor CXCR4, F2R, FLT1 (VEGFR1), KDR (VEGFR2), MET, IGF2R, LDLR and TFRC, and Carom endocytic partners GRASP, ITSN1-2, UBC, VCP and WASL are highly expressed. Carom is highly expressed in EC, LYM, MC, Mφ and myoblast. Abbreviation: CMC, Cardiomyocyte; VSMC, vascular smooth muscle cell; EC, endothelial cell; LYM, lymphocyte; MC, Monocyte; Mφ, Macrophage; GPCR, G-protein coupled receptor; RTK, Receptor tyrosine kinase; TRM, transmembrane receptor; other refer to table 1 and 3.




Figure 4. Models of Carom, endocytic partner and membrane receptor complexes. Interaction of membrane receptors with Carom and Carom partners were identified via NCBI and String databases. Solid lines indicate known interaction deposited in NCBI Gene database, which was established from affinity capture-MS, affinity capture-RNA, affinity capture-western, reconstituted complex and two-hybrid experimental data. Dashed lines indicate computational-predicted interaction in String database based on analyzing genomic information ('genomic context'-methods) or from transferring associations/interactions between organisms ('interolog'-transfer). Letters in red indicate genes with comparable high levels of expression paralleled with high levels of Carom in human cells identified in figure 3B. Noted that Carom may directly bind to TGBR1 (model A), indirectly interact with EGFR, ERBB2, ERBB3 and ERBB4 via Carom partners DAPP1, ITSM1, ITSN2, WAS, and WASL (model B), bind to receptors through its partner UBC (model C), and directly interact with partners GRASP, UBD and VCP (model D, E and F). Abbreviation: refer to table 1 and 3.

family appeared to be internalized by clathrin/caveolae-independent endocytosis (Indt) (table 1) (24).

4.2. Key steps in vesicle formation in CDE and its regulating proteins.

There are four steps during CDE process: 1) curvature initiation, 2) vesicle formation, 3) vesicle scission, and 4) un-coating vesicle (Figure 2). A group of proteins are involved in regulating endocytic vesicle formation. At first, the F-BAR protein binds to the plasma membrane and initiates membrane curvature and clathrin-coated endocytic vesicle formation (25, 26). F-BAR protein also can recruit adaptor proteins via its SH3 domain during vesicle formation. Clathrin is translocated to the site of the adaptor-concentrated membrane from the cytosol via adaptor proteins (2). The adaptor proteins link membrane cargo to clathrin and accessory proteins to form clathrin-coated endocytic vesicle. GTPase dynamin can then bind to the membrane and trigger vesicle scission and release upon GTP hydrolysis. Finally, ATPase heat shock cognate 70 (HSC70) binds to clathrin, disassociates clathrin, intersectin and dynamin from the vesicle and produces an un-coated endocytic vesicle containing the cargo molecules (27) (Figure 2).

4.3. F-BAR protein participates in membrane receptor endocytosis

It is reported that F-BAR protein mediates membrane receptor endocytosis by initiating

membrane curvature and endocytic vesicle formation in clathrin/caveolae-dependent endocytosis (8). F-BAR protein has 9 family members based on domain characterization. F-BAR protein contains one F-bar domain which binds to the cell membrane and other domains which interact with other molecules, such as Src homology-3 (SH3), Src homology-2 (SH2), protein kinase C-related kinase homology region 1 (HR1), F-BAR extension (FX), tyrosine kinase (Tyr-kinase), asparagine proline phenylalanine (NPF motif), μ -homology domain (μ HD) and Rho GTPase-activating protein (RhoGAP) domains (Table 2). 6 out of 9 F-BAR subfamily proteins are involved in membrane receptor endocytosis. CIP4 subfamily F-BAR proteins are required for EGFR internalization, trafficking and degradation (28). FCHO subfamily proteins are involved in LDLR endocytosis (29). PACSIN subfamily proteins participate in EGFR and transferrin/TFRC complex endocytosis (30). PSTPIP subfamily proteins can suppress transferrin/TFRC complex endocytosis (31). NOSTRIN subfamily protein is involved in assembling NOSTRIN-FGFR1-Rac1-Sos1 complex and regulating FGF signaling (28). FCHSDs subfamily proteins promote F-actin polymerization and membrane curvature which are key early events of endocytosis. However, endocytic receptors interacting with FCHSD protein have not been identified.

4.4. Identification of 9 Carom endocytic partners

The FCHSD2 gene encodes a protein termed as Carom, which is a novel membrane-associated

Table 4. Subcellular localization

Protein	Main subcellular locations (confidential levels)	Co-localization site with Carom				Nuclear localization Signal
		NC	CSK	CP	PM	
A. Carom and endocytic partner						
Carom	NC (5), CSK (2), CP (2), PM (2)	√	√	√	√	√
DAPP1	NC (2), CP (5), PM (5)	√	√	N/A	√	√
ITSN1	NC (2), CSK (2), CP (5), PM (5), E (1)	√	√	√	√	√
ITSN2	NC (2), CSK (5), CP (2), PM (2), EX (5), Gol (1)	√	√	√	√	√
UBC	NC (5), CSK (1), CP (5), PM (4), E (4), EX (5),	√	√	√	√	N/A
WAS	NC (4), CSK (5), CP (5), PM (2), E (2), EX (5), Mit (1)	√	√	√	√	N/A
WASL	NC (5), CSK (5), CP (5), PM (5), E (1), EX (5), Gol (3), Mit (2)	√	√	√	√	√
B. Endocytic membrane receptor						
ADRB1	CP (1), PM (5), E (5), ER (1), EX (2), Gol (1)	N/A	N/A	√	√	√
ADRB2	NC (3), CSK (2), CP (1), PM (5), E (5), ER (2), Lys (5)	√	√	√	√	N/A
1. CXCR4	NC (2), CSK (2), CP (5), PM (5), E (5), Lys (5)	√	√	√	√	√
2. EGFR	NC (5), CSK (3), CP (5), PM (5), E (5), ER (5), Gol (5), Lys (3), Mit (2)	√	√	√	√	√
3. ERBB2	NC (5), CSK (2), CP (3), PM (5), E (5), Lys (2)	√	√	√	√	√
ERBB3	NC (2), CSK (3), CP (1), PM (5)	√	√	√	√	N/A
4. ERBB4	NC (5), CSK (1), CP (5), PM (5), E (1), Lys (1), Mit (5)	√	√	√	√	√
FGFR1	NC (5), CSK (2), CP (5), PM (5), E (1), ER (1), Lys (1)	√	√	√	√	N/A
FGFR2	NC (5), CSK (2), CP (1), PM (5), E (1), ER (2), Gol (5), Lys (1)	√	√	√	√	N/A
FGFR3	NC (3), CSK (2), PM (5), ER (5), Gol (5), Lys (3)	√	√	N/A	√	N/A
FGFR4	NC (5), CSK (2), CP (3), PM (5), E (5), ER (5), Gol (5)	√	√	√	√	N/A
5. FLT1 (VEGFR1)	NC (2), CSK (2), CP (2), PM (5), E (5), ER (2), EX (2)	√	√	√	√	√
FOLR1	NC (5), PM (5), E (3), ER (4), EX (5),Gol (4)	√	N/A	N/A	√	N/A
F2R	NC (1), CSK (2), CP (4), PM (5), E (5), Gol (4)	√	√	√	√	N/A
6. IGF1R	NC (2), CSK (2), CP (2), PM (5), E (1), ER (1), EX (3), Mit (2)	√	√	√	√	√
7. IGF2R	NC (4), CSK (1), CP (2), PM (5), E (5), EX (5), Gol (5), Lys (4), Mit (1)	√	√	√	√	√
IL2RB	CP (1), PM (5), ER (1), EX (2), Gol (1)	N/A	N/A	√	√	N/A
IL2RG	NC (1), CSK (1), CP (1), PM (5), EX (3), Lys (1)	√	√	√	√	√
8. KDR (VEGFR2)	NC (5), CSK (2), CP (2), PM (5), E (5), ER (5), Gol (5), Lys (1), Mit (1)	√	√	√	√	√
9. LDLR	NC (2), CSK (1), CP (2), PM (5), E (5), ER (2), Gol (5), Lys (5), Mit (1)	√	√	√	√	√
10. MET	NC (2), CSK (2), CP (3), PM (5), ER (2)	√	√	√	√	√
11. NTRK1	NC (5), CSK (5), CP (5), PM (5), E (5), Gol (5), Lys (2), Mit (2)	√	√	√	√	√
12. PDGFRA	NC (5), CSK (2), CP (3), PM (5), EX (1)	√	√	√	√	√
TFRC	NC (2), CSK (2), CP (3), PM (5), E (5), ER (2), Lys (2), Mit (3)	√	√	√	√	N/A
13. TGFB1	NC (2), CSK (1), CP (1), PM (5), E (2), ER (2)	√	√	√	√	√
TGFB2	NC (2), CSK (1), CP (5), PM (5), Mit (2)	√	√	√	√	N/A

Subcellular localization of Carom partner and membrane receptor, and co-localization with Carom (Compartments database/cNLS database). Subcellular localization of Carom, partner and membrane receptors which can form 3 Carom-receptor complex models in figure 4, were determined in Compartments database. A. Subcellular localization and co-localization of Carom with endocytic partner. B. Subcellular localization of membrane receptor and co-localization with Carom. Noted that most of the Carom partners and membrane receptors co-localize with Carom in PM, CP and NC. Some of the Carom partners and membrane receptors contain NLS. 13 receptors can be potentially trans-localized to the nucleus because of the recognized NC localization and NLS, and labeled with numbers. Bolded words refer to molecules co-localized with Carom in subcellular sites as indicated in A. Numbers in parenthesis are confidence scores provided by Compartments database. Co-localization site with Carom are determined by present within the same cellular compartment and indicated by check marks (√). Nuclear localization signal was identified by using cNLS Mapper. Abbreviation: CP, cytoplasmic; CSK, cytoskeleton; E, endosome; ER, endoplasmic reticulum; EX, extracellular, Gol, golgi apparatus; Lys, lysosome; Mit, mitochondrion; NC, nucleus; PM, plasma membrane, others refer to table 1 and 3.

Table 5. Analysis of Carom:partner-receptor complex intracellular trafficking and function

Carom-receptor complex models (Carom:Partner-receptor)	Predicted Carom complex trafficking				
	Only in PM	PM to CP		to NC	
		Function	PMID#	Function	PMID#
A. Carom-TGFBR1 complex		TGFBR1 signaling/ degradation	21295082	RNA processing	22473997
B. Carom:partner-ERBBs complex Carom:DAPP1-ERBB3		ERBB3 signaling/degradation	22436610	N/A	
Carom:ITSN1-ERBB2, ERBB4, EGFR		ERBB2/4, EGFR signaling/ degradation	23472148	Transcriptional regulation/ DNA repair	26719328
Carom:ITSN2-ERBB2, ERBB4, EGFR		ERBB2/4, EGFR signaling/ degradation	23472148	Transcriptional regulation/ DNA repair	20670598
Carom:WAS-EGFR		EGFR signaling/degradation	23472148	N/A	
Carom:WASL-EGFR		EGFR signaling/degradation	23472148	Transcriptional regulation/ DNA repair	22127113
C. Carom:UBC-receptor complex Carom:UBC-ADRB1,ADRB2, CXCR4, EGFR, ERBB3, ERBB4, FGFR1, FGFR2, FGFR3, FGFR4, FLT1, FOLTR1, F2R, IGF1R, IGF2R, IL2RB, IL2RG, KDR, LDLR, MET, NTRK, PDGFRA, TFRC, TGFBR1, TGFBR2	Carom:UBC- FOLTR1	Ubiquitination, degradation		N/A	

Analysis of Carom-receptor complex intracellular trafficking and function. Carom-receptor complex trafficking and signaling are analyzed using information presented in figure 4 (interaction) and table 4 (co-localization). Carom-receptor complex functions are predicted based on their binding partner and receptor signaling reported in the literature (PMID#). We characterized 3 different Carom-receptor complex intracellular trafficking patterns: 1) only in PM, 2) PM to CP/CSK and 3) PM to NC. In 3 Carom-receptor complex models, the Carom-TGFBR1 and Carom:partner-ERBBs complex can be transferred to CP and CSK via endocytosis for signaling transduction and lysosomal degradation. They can also be transported into the nucleus and participate in RNA processing, transcription regulation and DNA repair because of the recognized NC localization and the detected NLS, except for Carom:DAPP1-ERBB3 and Carom:WAS-EGFR complex. The Carom:UBC-receptor complexes involve a large group of 24 receptors for ubiquitination and proteasome degradation by UBC, which lacks of NLS. Abbreviation: refer to table 1, 3 and 4.

protein with unknown function. Similar to most of the F-BAR family proteins, Carom binds to the cell membrane via N-terminal F-BAR domain association with membrane phospholipids and bridges the membrane with cytoskeleton. It interacts with proline-rich proteins, such as adaptor and signaling proteins, via its C-terminal SH3 domains to form a functional complex at cellular membranes. Based on their endocytosis-related functions reported in the literature, we selected 9 proteins from 26 Carom partners we previously identified, and termed them as Carom endocytic partners (Table 3) (32). It was reported that these Carom endocytic partners played important role in regulating actin polymerization, endocytic vesicle formation and protein ubiquitination. The Carom endocytic partner proteins contain different domains, such as SH2, Pleckstrin homology (PH), Ubiquitin, CDC42-Rac interactive binding (CRIB), and WASP-Homology 1/2 (WH1/2) domain, which can recruit proteins to organize signaling complexes at cellular membranes. Carom partners can bind to receptors. We found that DAPP1 binds to ERBB3 and that ITSN2/UBC/WAS/WASL bind to EGFR from experimental data generated via affinity capture-MS, affinity capture-RNA, affinity capture-western, reconstituted complex and two-hybrid technologies, and deposited in NCBI database at the gene/interaction branch. It

is well documented that Carom endocytic partners ITSN1/UBC/UBD, VCP and WASL regulate EGFR ubiquitination, degradation and endocytosis process (8, 28, 29, 33, 34).

4.5. Endocytic membrane receptor, Carom and its endocytic partner are differently expressed in human primary cells

We anticipate that the expression of Carom, its corresponding partners and endocytic receptors is comparable in the cells where Carom-organized receptor endocytosis takes place, and examined their relevant expression in 12 human body systems and 10 primary cells (Figure 3). Gene expression levels (mRNA) were obtained from microarray database (<https://www.genevestigator.com/gv/>). We found that human body systems do not have distinguished patterns of Carom, endocytic membrane receptors and partners' expression. GPCR class endocytic membrane receptors had relatively low levels of expression in most of the body system, but were highly expressed in immune system and in monocyte (MC), macrophage (M ϕ) and pancreatic islet cells. A few of RTK class receptors, such as EGFR, KDR (VEGFR2), MET, PDGFRA, appeared to be highly expressed in most of the body systems and in circulatory system

cells, including cardiomyocyte (CMC), aortic vascular smooth muscle cell (VSMC), and aortic endothelial cell (EC). Cytokines and growth factor-related TRM class receptors had low levels of expression in most of body systems, but highly expressed in immune system (MC & M ϕ) and pancreatic islet cells. In contrast, TRM class receptors LDLR and TFRC, which transport lipid and iron into the cells, were highly expressed in all human body systems and cells. Carom and its partners were expressed at medium levels in all body systems. In the circulatory system cells (Figure 3), Carom was highly expressed in EC paralleled with highly expressed receptors (CXCR4, F2R, FLT1 (VEGFR1), KDR (VEGFR2), MET, IGF2R, LDLR and TFRC) and partners (GRASP, ITSN1-2, UBC, VCP and WASL). In lymphocyte, Carom was highly expressed paralleled with highly expressed receptors (EGFR, ERBB3, MET, IGF1-2R, LDLR and TFRC) and partners (DAPP1, ITSN2 and WAS). In MC, highly expressed Carom was paralleled with receptors (CCR5, CXCR4, CSF1R, TGFBR1, IGF2R and FOLR3) and partners (DAPP1, ITSN2 and WAS). Macrophage had higher levels of Carom expression paralleled with a large group of highly expressed receptors (CCR5, CXCR4, CSF1R, FLT1 (VEGFR1), TGFBR1, IGF2R, FOLR3 and TFRC) and partners (DAPP1, ITSN1-2, UBC, UBD, WAS and WASL), which is comparable with that in EC.

4.6. Carom may directly or indirectly bind to endocytic membrane receptor through partner protein

To generate models of Carom-receptor complexes for receptor endocytosis and signaling, we analyzed the interaction of Carom with membrane receptors and partners using information from NCBI experimental database and computational String database (Figure 4). We proposed 3 Carom-receptor complex models: A) Carom-TGFBR1 complex, in which Carom binds directly to TGFBR1, B) Carom:partner-ERBBs complex, in which Carom indirectly interacts with ERBB2, ERBB3, ERBB4, EGFR via Carom partners DAPP1, ITSN1, ITSN2, WAS, and WASL, C) Carom:UBC-receptor complex in which Carom may bind to 25 of 34 membrane receptors (ADRB1, ADRB2, CXCR4, EGFR, ERBB3, ERBB4, FGFR1, FGFR2, FGFR3, FGFR4, FLT1 (VEGFR1), FOLTR1, F2R, IGF1R, IGF2R, IL2RB, IL2RG, KDR (VEGFR2), LDLR, MET, NTRK1, PDGFRA, TFRC, TGFBR1, TGFBR2) through partner UBC. In addition, Carom directly interacts with partners GRASP, UBD and VCP (model D, E and F) and this aids in the transduction of their signals.

4.7. Carom co-localization with endocytic membrane receptors and partners in different cellular micro-compartment

Interacting complexes are more likely to be presented within the same cellular compartment. We analyzed subcellular localization of Carom-

receptor complex proteins identified in Figure 4 using Compartments database from manually curated literature, high-throughput screens, automatic text mining, and sequence-based prediction methods. We found that Carom is located in 4 major cell compartments, PM, NC, CP and CSK with the highest confidential level in the NC (confidential level 5) (Table 4). Carom endocytic partners (DAPP1, ITSN1/2, UBC, WAS and WASL) are distributed in multiple compartments and mostly co-localized with Carom in 4 major cell compartments, with the exception of DAPP1 which is not located in the CSK. Except for UBC and WAS, all Carom-receptor complex related partner proteins contain NLS. Carom-related membrane receptors are localized in multiple cell compartments (Table 4). While looking at the 4 Carom-existent cell compartments, most of the membrane receptors can co-localize with Carom, except that ADRB1 was only located on the PM and CP, and FOLR1 is not sited in CP and CSK. Interestingly, FGFR3 is located in the PM, CSK and NC, but not in CP. We identified 13 receptors which can be potentially trans-localized to the nucleus because of the recognized NC localization and the detected NLS.

4.8. Carom-receptor complex intracellular trafficking and function

We analyzed intracellular trafficking and function of Carom-receptor complexes identified in Figure 4. Three models of Carom-receptor complexes are listed as, A) Carom-TGFBR1, B) Carom:partner-ERBBs, and C) Carom:UBC-receptor (Table 5). Based on their co-localization in the subcellular compartment and NLS, we characterized 3 different Carom-receptor complex intracellular trafficking patterns: 1) only in PM, 2) PM to CP/CSK, and 3) PM to NC. Among 13 potential nuclear trans-localized receptors identified, 9 receptors were found to interact with Carom via UBC which does not contain NLS. These 9 receptors were not justified as nuclear trans-localized receptors. The function of Carom-receptor complexes was determined based on signaling information related with its binding partner or receptor in literature.

As summarized in Table 5, the Carom-TGFBR1 complex can be transferred to CP and CSK via endocytosis for signaling transduction and lysosomal degradation (35). It can also be transported into the nucleus and participate in RNA processing (36). Carom:partner-ERBBs complex can be transferred to CP and CSK for signaling transduction and lysosomal degradation (37). Most of the Carom:partner-ERBBs complexes, except for Carom:DAPP1-ERBB3 and Carom:WAS-EGFR, can be transported into the nucleus to regulate transcription and DNA repair (38). The Carom:UBC-receptor complexes involve a large group of 24 receptors, including receptors for cytokine, growth factor, that are directed by UBC which lacks

of NLS, and are subjected to ubiquitination and proteasome degradation (37, 39, 40). The Carom:UBC-FOLR1 complex may only stay in PM, because FOLR1 does not exist in CP and CSK.

5. DISCUSSION

Endocytic trafficking of molecules is a highly regulated process involving multiple steps and molecules (Figure 1&2). In response to ligands stimulation, the BAR super family proteins can bind to the cell membrane and bend to either positive or negative curvature. BAR proteins then recruit other adaptor proteins or accessory proteins to the deformed membrane to form endocytic vesicles. After endocytosis, cargos are destined to different subcellular organelles, including different endosomes and lysosomes.

Receptor trafficking is an important pathway for their signaling. The previous concept that receptor endocytosis would only contribute to its signal attenuation has already been challenged. Recent evidence demonstrated that receptor endocytosis and the following subcellular organelle redistribution regulate downstream signaling and gene regulation (41-44). As summarized in table 1, at least 34 membrane receptors can be internalized via mostly CDE or CIE mediated endocytosis process, which contribute to their functions of regulating cell differentiation, proliferation, survival, angiogenesis, tumor transformation and immune regulation.

Recently, F-BAR proteins, a subfamily of the BAR superfamily, have been identified as important coordinators that regulate endocytosis. In general, F-BAR proteins bind to the cell membrane via the association of F-BAR domain with membrane phospholipids. Through the SH3 domain, F-BAR proteins interact with WASP or GTPase dynamin to regulate the initiation and scission of the endocytic vesicle. We found out that at least 4 F-BAR protein subfamilies (CIP4, FCHO, PACSIN and NOSTRIN) are involved in the formation of endocytic vesicles and the assembly of endocytic complexes (Table 2) (7, 8). We listed 4 receptor endocytosis mechanisms, including CIP4 subfamily-related EGFR degradation, NOSTRIN subfamily-regulated FGFR signaling, FCHO2-regulated LDLR endocytosis and PACSIN3/PSTPIP1-regulated TFRC endocytosis. These findings presented fundamental mechanisms for F-BAR protein-mediated receptor endocytosis. F-BAR protein-mediated receptor endocytosis, although less studied, may play critical roles in growth control, angiogenesis and lipid metabolism.

The FCHSD subfamily has two members, FCHSD1 and FCHSD2, each containing of one F-BAR domain and two SH3 domains. The biological function

of FCHSD subfamily proteins may be related to F-actin polymerization based on their direct interaction with WASP in *E. Coli* to promote WASP-Arp2/3-dependent F-actin polymerization (7). WASP is known to bind to Arp2/3 complex, via its C-terminal, to nucleate actin filaments, which then elongate at their free barbed ends to induce F-actin polymerization (45, 46). FCHSD2 (Carom) is a newly identified FCHSD subfamily member with unknown function. It is suggested that acute myeloid leukemia (AML) patients with high Carom expression have increased leukemia chemoresistance. We have previously proposed that Carom may regulate membrane curvature, promote F-actin polymerization and recruit adaptor proteins via its partner in the process of CDE (25). Carom related membrane receptor and endocytosis have not been studied. We hypothesized that Carom may regulate receptor endocytosis via its partner proteins and identified 9 Carom endocytic partners (Table 3). We found that Carom partners DAPP1 can bind to ERBB3 and that ITSN2/UBC/WAS/WASL can bind to EGFR. ITSN1/UBC/UBD/VCP bind to both Carom and EGFR leading to EGFR ubiquitination and degradation (47). We hypothesize that Carom regulates EGFR and other receptor internalization and signaling via interaction with its endocytic partners.

In the efforts to explore the functional connection of Carom and related receptors, we examined cell type expression profile of Carom, its endocytic binding partners and membrane receptors in human body systems and primary cells (Figure 3). High level Carom expression was found paralleled with some highly expressed endocytosis-related membrane receptors and Carom partners in aortic endothelial cell, lymphocytes, monocytes and macrophages, which usually display robust endocytosis phenomena. These results indicate that Carom and its partners regulate endocytosis-related endothelial function and myeloid cell related innate immune function.

We further analyzed the subcellular localization of these proteins to search potential signal partners in cell organelles in Carom-related membrane receptors endocytosis (Table 4). We found that Carom is located in all major subcellular domains, including NC, CSK, CP and PM, which is a typical pattern of trafficking signal molecules. The co-localization relationship of Carom with different endocytic binding partners and membrane receptors is dynamic. It appears that Carom co-localizes with all membrane receptors and endocytic binding partners at plasma membrane, suggesting the critical role of Carom in the initial step of receptor endocytosis on plasma membrane. In the NC, Carom is co-localized with all partners and most of the receptors in the NC, except for ADRB1 and IL2RB. We identified NLS in Carom and hypothesized that Carom can be translocated into the nucleus and is responsible for taking the




Figure 5. Hypothetic working model of Carom-mediated membrane receptor trafficking and endocytosis. Carom can form 3 types of receptor complexes and mediate membrane receptor trafficking and endocytosis. A) Carom-TGFBR1 complex can enter the nucleus via nuclear localization signal and facilitate RNA processing. B) Carom:Partner-ERBBs complexes that Carom associates with receptor via its partner, such as Carom:DAPP1-ERBB3; Carom:ITSN1-ERBB2, ERBB4, EGFR; Carom:ITSN2-ERBB2, ERBB4, EGFR; Carom:WAS-EGFR; Carom:WASL-EGFR (details in Table 5). Carom:Partner-ERBBs complex (Carom:ITSN1-ERBB2, ERBB4, EGFR; Carom:ITSN2-ERBB2, ERBB4, EGFR; Carom:WASL-EGFR) can be transported into the nucleus, bind to transcriptional factor and promote transcription, or bind to damaged DNA to facilitate DNA repair via activation of DNA-PK. Both Carom-TGFBR1 and Carom:Partner-ERBBs complexes can facilitate receptor signaling in the cytoplasm leading to proliferation, differentiation and tumor transformation, and can be degraded in lysosome. C) Carom:UBC-Receptor complexes may bind to its partner UBC, which further bind to 24 membrane receptors and facilitate ubiquitination and proteasome degradation. Abbreviation: DNA-PK, DNA-protein kinase; others refer to table 1 and 3.

Carom:partner-receptor complexes into the nucleus, because the NC-localized receptors (FGFR1-4, ERBB3, FOLR1, F2R, TFRC, TGFBR2) and partners (UBC and WAS) do not have an identified NLS. We found that except for FGFR3 and FOLR1, Carom and most of the receptors and partners are also located in the CP, suggesting that a proportion of Carom:partner-receptor complexes can be disassociated from membrane structure and organelles, and released to the CP. The dynamic distribution of the component of Carom:partner-receptor complexes in various subcellular domain and organelles, including CSK, endosome (E), endoplasmic reticulum (ER), lysosome (Lys), mitochondrion (Mit) presented different intracellular trafficking pathways for Carom:partner-receptor complexes from CP to the NC.

Based on above findings, we presented three novel models for Carom-related receptor trafficking (Table 5 & Figure 5); Carom can A) directly bind to receptor (TGFBR1), B) indirectly binds to receptor through its partner (EGFR, ERBB2-4 receptors) to initiate the formation of endocytic vesicle, and C) facilitate membrane endocytosis through ubiquitination related proteins (UBC). Model C is likely responsible for endocytosis, ubiquitination and proteasome degradation of a large group of receptors. It is noticed that ubiquitin itself is a sorting signal for membrane receptor endocytosis. There exist different sorting

machineries that determine how receptors are selected by compartment specific ubiquitin-binding proteins and are delivered to cellular destination (48, 49). These three models suggest that Carom may play a critical role in regulating intracellular trafficking and signaling of a large number of membrane receptors.

Transforming growth factor β (TGF- β) plays a critical role in embryogenesis and adult tissue development by regulating cell proliferation, differentiation, and migration (50). It is suggested that TGFBR1 plays an important regulatory role in TGF- β signaling via CDE to promote TGF- β -induced Smad activation and transcriptional regulation or Caveolae-dependent endocytosis to facilitate the degradation of TGF- β (35). We proposed a novel Carom-TGFBR1 complex for TGF- β signaling through database mining (Table 5). ERBBs family contains EGFR, ERBB2, ERBB3 and ERBB4. It is suggested that ERBBs are internalized mainly through CDE after ligand binding, which is followed by receptor activation or lysosomal degradation (51-53). Our data suggests that Carom can bind to ERBBs via its partners DAPP1, ITSN1/2, WAS and WASL, and such interaction complexes contribute to receptor endocytosis and control of signaling (Table 5). UBC gene encodes polyubiquitin-C protein, which is involved in the regulation of CDE and protein ubiquitination (54-56). Through our analysis, we anticipate that Carom may mediate those 24

membrane receptors transport into cytoplasm and cytoskeleton via interacting with UBC for ubiquitination. As mentioned above, Carom may regulate receptor endocytosis through specific sorting machinery for individual receptor.

It is suggested that certain endocytic proteins translocate to the nucleus in response to extracellular signals, which may affect gene transcription and chromatin remodeling machinery (57-59). The mechanism by which endocytic proteins enter the nucleus is based on NLS or interaction with carrier proteins (21). (Table 4). Carom and its endocytic partners (DAPP1, ITSN1-2 and WASL) were identified to have NLS through protein sequence analysis. Therefore, we propose that Carom-TGFBR1 and Carom:partner-ERBBs complexes may transport from the membrane to the nucleus to activate TGFBR1 and ERBB signaling for RNA processing, gene transcription or DNA repair (36, 60) (Table 5).

Traditional concept recognizes that the purpose of endocytosis of membrane receptors is to terminate receptor mediated signaling. However, it is now recognized that receptor internalization, especially for RTK families, is highly regulated via various mechanisms. For example, EGFR and FGFR employ different molecular mechanisms for nuclear translocation (60, 61). Unlike EGFR which displays NLS, FGFR is translocated into the nucleus from early endosome (62). EGFR endocytosis is required for optimal activation of sub-populations of signal transducers (63). EGFR endocytosis and post-endocytic traffic display versatile pathways and such traffic can lead to different cellular behaviors, such as proliferation, survival, tumorigenesis and DNA repair (Figure 5). Different KDR (VEGFR2) trafficking pathways via different subcellular compartments affect different cellular behaviors, ranging from proliferation, migration, tumorigenesis and blood vessel formation (42, 64). The proposed 3 models in figure 5 presented a simple network. Studies to define the subcellular localization of Carom:partner-receptor complexes in early and late endosome, lysosome or trans Golgi, nucleus should provide strong evidence and discover relevant molecular mechanisms.

A more complex regulatory network could be involved in regulating Carom-mediated receptor endocytosis and trafficking to the nucleus, especially considering the two SH3 domains which can associate with many other adaptor proteins. SH3 signaling may lead to phosphorylation of the receptor and adaptor proteins. The detailed mechanisms of surface receptors translocation to the nucleus are largely unveiled. Whether the nuclear translocated receptors come from receptors embedded in endosomes are under debate and requires experimental clarification (61, 65).

The combination of various bioinformatics tools employed in the study is very powerful for the identification of protein complexes involving complicated intracellular trafficking mechanism. While more and more online large databases become available, it is possible to develop model systems regulating important biological process and to predict molecular targets. The identified mechanistic model system can be important guidance for future experimental science and may lead to the discovery of a novel mechanism for human diseases and therapeutic targets.

6. CONCLUSION

In this study, we identified 34 endocytic membrane receptors and 9 Carom endocytic partners and established their expression profiles in human primary cells. We established 3 models of Carom-receptor complexes and their intracellular trafficking based on protein-protein interaction and subcellular localization. We proposed that F-BAR protein Carom may mediate receptor endocytosis and transport endocytic receptors to the cytoplasm for receptor signaling and lysosome/proteasome degradation, or to the nucleus for RNA processing, gene transcription and DNA repair.

7. ACKNOWLEDGMENTS

This work was supported in part by National Institutes of Health (NIH) Grants number: HL67033, HL77288, HL82774, HL110764 and HL117654 (HW); HL9445, HL108910 and HL116917 (XFY). The authors declare that they have no conflict of interest.

8. REFERENCES

1. A. Du Toit: Endocytosis. A new gateway into cells. *Nat Rev Mol Cell Biol*, 16(2), 68 (2015)
DOI: 10.1038/nrmicro3425
DOI: 10.1038/nrm3946
DOI: 10.1038/nrm3939
DOI: 10.1038/nrmicro3426
PMid:25604192
2. H. T. McMahon and E. Boucrot: Molecular mechanism and physiological functions of clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol*, 12(8), 517-33 (2011)
DOI: 10.1038/nrm3151
PMid:21779028
3. Z. J. Cheng, R. D. Singh, D. L. Marks and R. E. Pagano: Membrane microdomains, caveolae, and caveolar endocytosis of sphingolipids. *Mol Membr Biol*, 23(1), 101-10 (2006)
DOI: 10.1080/09687860500460041
PMid:16611585

4. R. C. Aguilar and B. Wendland: Endocytosis of membrane receptors: two pathways are better than one. *Proc Natl Acad Sci U S A*, 102(8), 2679-80 (2005)
DOI: 10.1073/pnas.0500213102
PMid:15710869 PMCID:PMC549498
5. G. Kourouniotis, Y. Wang, S. Pennock, X. Chen and Z. Wang: Non-Ligand-Induced Dimerization is Sufficient to Initiate the Signalling and Endocytosis of EGF Receptor. *Int J Mol Sci*, 17(8) (2016)
DOI: 10.3390/ijms17081200
PMid:27463710 PMCID:PMC5000598
6. D. P. Poole and N. W. Bunnett: G Protein-Coupled Receptor Trafficking and Signalling in the Enteric Nervous System: The Past, Present and Future. *Adv Exp Med Biol*, 891, 145-52 (2016)
DOI: 10.1007/978-3-319-27592-5_14
PMid:27379642
7. H. Cao, X. Yin, Y. Cao, Y. Jin, S. Wang, Y. Kong, Y. Chen, J. Gao, S. Heller and Z. Xu: FCHSD1 and FCHSD2 are expressed in hair cell stereocilia and cuticular plate and regulate actin polymerization *in vitro*. *PLoS One*, 8(2), e56516 (2013)
DOI: 10.1371/journal.pone.0056516
PMid:23437151 PMCID:PMC3577914
8. W. M. Henne, E. Boucrot, M. Meinecke, E. Evergren, Y. Vallis, R. Mittal and H. T. McMahon: FCHO proteins are nucleators of clathrin-mediated endocytosis. *Science*, 328(5983), 1281-4 (2010)
DOI: 10.1126/science.1188462
PMid:20448150 PMCID:PMC2883440
9. X. Huang, R. Gong, X. Li, A. Virtue, F. Yang, I. H. Yang, A. H. Tran, X. F. Yang and H. Wang: Identification of novel pretranslational regulatory mechanisms for NF-kappaB activation. *J Biol Chem*, 288(22), 15628-40 (2013)
DOI: 10.1074/jbc.M113.460626
PMid:23515310 PMCID:PMC3668723
10. N. C. Chen, F. Yang, L. M. Capecchi, Z. Gu, A. I. Schafer, W. Durante, X. F. Yang and H. Wang: Regulation of homocysteine metabolism and methylation in human and mouse tissues. *FASEB J*, 24(8), 2804-17 (2010)
DOI: 10.1096/fj.09-143651
PMid:20305127 PMCID:PMC2909276
11. X. Li, J. Mai, A. Virtue, Y. Yin, R. Gong, X. Sha, S. Gutchigian, A. Frisch, I. Hodge, X. Jiang, H. Wang and X. F. Yang: IL-35 is a novel responsive anti-inflammatory cytokine--a new system of categorizing anti-inflammatory cytokines. *PLoS One*, 7(3), e33628 (2012)
DOI: 10.1371/journal.pone.0033628
PMid:22438968 PMCID:PMC3306427
12. H. Ilioka and I. G. Macara: Detection of RNA-Protein Interactions Using Tethered RNA Affinity Capture. *Methods Mol Biol*, 1316, 67-73 (2015)
DOI: 10.1007/978-1-4939-2730-2_6
PMid:25967053
13. M. Barrios-Rodiles, K. R. Brown, B. Ozdamar, R. Bose, Z. Liu, R. S. Donovan, F. Shinjo, Y. Liu, J. Dembowy, I. W. Taylor, V. Luga, N. Przulj, M. Robinson, H. Suzuki, Y. Hayashizaki, I. Jurisica and J. L. Wrana: High-throughput mapping of a dynamic signaling network in mammalian cells. *Science*, 307(5715), 1621-5 (2005)
DOI: 10.1126/science.1105776
PMid:15761153
14. M. Kotlyar, C. Pastrello, F. Pivetta, A. Lo Sardo, C. Cumbaa, H. Li, T. Naranian, Y. Niu, Z. Ding, F. Vafaei, F. Broackes-Carter, J. Petschnigg, G. B. Mills, A. Jurisicova, I. Stagljar, R. Maestro and I. Jurisica: In silico prediction of physical protein interactions and characterization of interactome orphans. *Nat Methods*, 12(1), 79-84 (2015)
DOI: 10.1038/nmeth.3178
PMid:25402006
15. Z. Lu and T. Hunter: Degradation of activated protein kinases by ubiquitination. *Annu Rev Biochem*, 78, 435-75 (2009)
DOI: 10.1146/annurev.biochem.013008.092711
PMid:19489726 PMCID:PMC2776765
16. M. Galovic, D. Xu, L. B. Areces, R. van der Kammen and M. Innocenti: Interplay between N-WASP and CK2 optimizes clathrin-mediated endocytosis of EGFR. *J Cell Sci*, 124(Pt 12), 2001-12 (2011)
DOI: 10.1242/jcs.081182
PMid:21610097
17. R. B. Jones, A. Gordus, J. A. Krall and G. MacBeath: A quantitative protein interaction network for the ErbB receptors using protein microarrays. *Nature*, 439(7073), 168-74 (2006)
DOI: 10.1038/nature04177
PMid:16273093

18. J. Tong, P. Taylor, S. M. Peterman, A. Prakash and M. F. Moran: Epidermal growth factor receptor phosphorylation sites Ser991 and Tyr998 are implicated in the regulation of receptor endocytosis and phosphorylations at Ser1039 and Thr1041. *Mol Cell Proteomics*, 8(9), 2131-44 (2009)
DOI: 10.1074/mcp.M900148-MCP200
PMid:19531499 PMCID:PMC2742444
19. F. Bornke: Corrigendum: The complex becomes more complex: protein-protein interactions of SnRK1 with DUF581 family proteins provide a framework for cell- and stimulus type-specific SnRK1 signaling in plants. *Front Plant Sci*, 5, 693 (2014)
DOI: 10.3389/fpls.2014.00054
20. X. Tang, X. Hu, X. Yang, Y. Fan, Y. Li, W. Hu, Y. Liao, M. C. Zheng, W. Peng and L. Gao: Predicting diabetes mellitus genes via protein-protein interaction and protein subcellular localization information. *BMC Genomics*, 17 Suppl 4, 433 (2016)
DOI: 10.1186/s12864-016-2795-y
PMid:27535125 PMCID:PMC5001230
21. B. Pyrzynska, I. Pilecka and M. Miaczynska: Endocytic proteins in the regulation of nuclear signaling, transcription and tumorigenesis. *Mol Oncol*, 3(4), 321-38 (2009)
DOI: 10.1016/j.molonc.2009.06.001
PMid:19577966
22. Y. Mosesson, K. Shtiegman, M. Katz, Y. Zwang, G. Vereb, J. Szollosi and Y. Yarden: Endocytosis of receptor tyrosine kinases is driven by monoubiquitylation, not polyubiquitylation. *J Biol Chem*, 278(24), 21323-6 (2003)
DOI: 10.1074/jbc.C300096200
PMid:12719435
23. C. Le Roy and J. L. Wrana: Clathrin- and non-clathrin-mediated endocytic regulation of cell signalling. *Nat Rev Mol Cell Biol*, 6(2), 112-26 (2005)
DOI: 10.1038/nrm1571
PMid:15687999
24. C. Basquin, M. Trichet, H. Vihinen, V. Malarde, T. Lagache, L. Ripoll, E. Jokitalo, J. C. Olivo-Marin, A. Gautreau and N. Sauvonnet: Membrane protrusion powers clathrin-independent endocytosis of interleukin-2 receptor. *EMBO J*, 34(16), 2147-61 (2015)
DOI: 10.15252/embj.201490788
PMid:26124312 PMCID:PMC4557667
25. S. Liu, X. Xiong, X. Zhao, X. Yang and H. Wang: F-BAR family proteins, emerging regulators for cell membrane dynamic changes-from structure to human diseases. *J Hematol Oncol*, 8, 47 (2015)
DOI: 10.1186/s13045-015-0144-2
PMid:25956236 PMCID:PMC4437251
26. I. K. Jarsch, F. Daste and J. L. Gallop: Membrane curvature in cell biology: An integration of molecular mechanisms. *J Cell Biol*, 214(4), 375-87 (2016)
DOI: 10.1083/jcb.201604003
PMid:27528656
27. R. Sousa, H. S. Liao, J. Cuellar, S. Jin, J. M. Valpuesta, A. J. Jin and E. M. Lafer: Clathrin-coat disassembly illuminates the mechanisms of Hsp70 force generation. *Nat Struct Mol Biol*, 23(9), 821-9 (2016)
DOI: 10.1038/nsmb.3272
PMid:27478930
28. J. Hu, F. Troglio, A. Mukhopadhyay, S. Everingham, E. Kwok, G. Scita and A. W. Craig: F-BAR-containing adaptor CIP4 localizes to early endosomes and regulates Epidermal Growth Factor Receptor trafficking and downregulation. *Cell Signal*, 21(11), 1686-97 (2009)
DOI: 10.1016/j.cellsig.2009.07.007
PMid:19632321
29. E. E. Mulkearns and J. A. Cooper: FCH domain only-2 organizes clathrin-coated structures and interacts with Disabled-2 for low-density lipoprotein receptor endocytosis. *Mol Biol Cell*, 23(7), 1330-42 (2012)
DOI: 10.1091/mbc.E11-09-0812
PMid:22323290 PMCID:PMC3315808
30. B. J. de Kreuk, E. C. Anthony, D. Geerts and P. L. Hordijk: The F-BAR protein PACSIN2 regulates epidermal growth factor receptor internalization. *J Biol Chem*, 287(52), 43438-53 (2012)
DOI: 10.1074/jbc.M112.391078
PMid:23129763 PMCID:PMC3527931
31. K. M. Cooper, D. A. Bennin and A. Huttenlocher: The PCH family member proline-serine-threonine phosphatase-interacting protein 1 targets to the leukocyte uropod and regulates directed cell migration. *Mol Biol Cell*, 19(8), 3180-91 (2008)
DOI: 10.1091/mbc.E08-02-0225
PMid:18480402 PMCID:PMC2488309

32. S. Liu, X. Xiong, S. V. Thomas, Y. Xu, X. Cheng, X. Zhao, X. Yang and H. Wang: Analysis for Carom complex, signaling and function by database mining. *Front Biosci (Landmark Ed)*, 21, 856-72 (2016)
DOI: 10.2741/4424
PMid:22520625 PMCID:PMC3418567
33. I. Kovacevic, J. Hu, A. Siehoff-Icking, N. Opitz, A. Griffin, A. C. Perkins, A. L. Munn, W. Muller-Esterl, R. Popp, I. Fleming, B. Jungblut, M. Hoffmeister and S. Oess: The F-BAR protein NOSTRIN participates in FGF signal transduction and vascular development. *EMBO J*, 31(15), 3309-22 (2012)
DOI: 10.1038/emboj.2012.176
PMid:22751148 PMCID:PMC3411079
34. J. Modregger, B. Ritter, B. Witter, M. Paulsson and M. Plomann: All three PACSIN isoforms bind to endocytic proteins and inhibit endocytosis. *J Cell Sci*, 113 Pt 24, 4511-21 (2000)
PMid:11082044
35. A. A. Bizet, K. Liu, N. Tran-Khanh, A. Saksena, J. Vorstenbosch, K. W. Finnsen, M. D. Buschmann and A. Philip: The TGF-beta co-receptor, CD109, promotes internalization and degradation of TGF-beta receptors. *Biochim Biophys Acta*, 1813(5), 742-53 (2011)
DOI: 10.1016/j.bbamcr.2011.01.028
PMid:21295082
36. M. Chandra, S. Zang, H. Li, L. J. Zimmerman, J. Champer, A. Tsuyada, A. Chow, W. Zhou, Y. Yu, H. Gao, X. Ren, R. J. Lin and S. E. Wang: Nuclear translocation of type I transforming growth factor beta receptor confers a novel function in RNA processing. *Mol Cell Biol*, 32(12), 2183-95 (2012)
DOI: 10.1128/MCB.00320-12
PMid:22473997 PMCID:PMC3372271
37. H. A. Alwan, E. J. van Zoelen and J. E. van Leeuwen: Ligand-induced lysosomal epidermal growth factor receptor (EGFR) degradation is preceded by proteasome-dependent EGFR de-ubiquitination. *J Biol Chem*, 278(37), 35781-90 (2003)
DOI: 10.1074/jbc.M301326200
PMid:12829707
38. Y. N. Wang and M. C. Hung: Nuclear functions and subcellular trafficking mechanisms of the epidermal growth factor receptor family. *Cell Biosci*, 2(1), 13 (2012)
DOI: 10.1186/2045-3701-2-13
PMid:22520625 PMCID:PMC3418567
39. G. Levkowitz, H. Waterman, E. Zamir, Z. Kam, S. Oved, W. Y. Langdon, L. Beguinot, B. Geiger and Y. Yarden: c-Cbl/Sli-1 regulates endocytic sorting and ubiquitination of the epidermal growth factor receptor. *Genes Dev*, 12(23), 3663-74 (1998)
DOI: 10.1101/gad.12.23.3663
PMid:9851973 PMCID:PMC317257
40. X. Long and K. P. Nephew: Fulvestrant (ICI 182,780)-dependent interacting proteins mediate immobilization and degradation of estrogen receptor-alpha. *J Biol Chem*, 281(14), 9607-15 (2006)
DOI: 10.1074/jbc.M510809200
PMid:16459337
41. A. Tomas, C. E. Futter and E. R. Eden: EGF receptor trafficking: consequences for signaling and cancer. *Trends Cell Biol*, 24(1), 26-34 (2014)
DOI: 10.1016/j.tcb.2013.11.002
PMid:24295852 PMCID:PMC3884125
42. A. Horowitz and H. R. Seerapu: Regulation of VEGF signaling by membrane traffic. *Cell Signal*, 24(9), 1810-20 (2012)
DOI: 10.1016/j.cellsig.2012.05.007
PMid:22617029
43. P. Balogh, S. Katz and A. L. Kiss: The role of endocytic pathways in TGF-beta signaling. *Pathol Oncol Res*, 19(2), 141-8 (2013)
DOI: 10.1007/s12253-012-9595-8
PMid:23274761
44. L. K. Goh and A. Sorkin: Endocytosis of receptor tyrosine kinases. *Cold Spring Harb Perspect Biol*, 5(5), a017459 (2013)
DOI: 10.1101/cshperspect.a017459
PMid:23637288 PMCID:PMC3632065
45. W. L. Lee, M. Bezanilla and T. D. Pollard: Fission yeast myosin-I, Myo1p, stimulates actin assembly by Arp2/3 complex and shares functions with WASp. *J Cell Biol*, 151(4), 789-800 (2000)
DOI: 10.1083/jcb.151.4.789
PMid:11076964 PMCID:PMC2169449
46. S. B. Padrick, L. K. Doolittle, C. A. Brautigam, D. S. King and M. K. Rosen: Arp2/3 complex is bound and activated by two WASP proteins. *Proc Natl Acad Sci U S A*, 108(33), E472-9 (2011)
DOI: 10.1073/pnas.1100236108
PMid:21676863 PMCID:PMC3158169

47. M. N. Okur, J. Ooi, C. W. Fong, N. Martinez, C. Garcia-Dominguez, J. M. Rojas, G. Guy and J. P. O'Bryan: Intersectin 1 enhances Cbl ubiquitylation of epidermal growth factor receptor through regulation of Sprouty2-Cbl interaction. *Mol Cell Biol*, 32(4), 817-25 (2012)
DOI: 10.1128/MCB.05647-11
PMid:22158968 PMCID:PMC3272971
48. R. C. Piper, I. Dikic and G. L. Lukacs: Ubiquitin-dependent sorting in endocytosis. *Cold Spring Harb Perspect Biol*, 6(1) (2014)
DOI: 10.1101/cshperspect.a016808
PMid:24384571 PMCID:PMC3941215
49. K. Haglund and I. Dikic: The role of ubiquitylation in receptor endocytosis and endosomal sorting. *J Cell Sci*, 125(Pt 2), 265-75 (2012)
DOI: 10.1242/jcs.091280
PMid:22357968
50. C. A. Clement, K. D. Ajbro, K. Koefoed, M. L. Vestergaard, I. R. Veland, M. P. Henriques de Jesus, L. B. Pedersen, A. Benmerah, C. Y. Andersen, L. A. Larsen and S. T. Christensen: TGF-beta signaling is associated with endocytosis at the pocket region of the primary cilium. *Cell Rep*, 3(6), 1806-14 (2013)
DOI: 10.1016/j.celrep.2013.05.020
PMid:23746451
51. M. M. Sak, K. Breen, S. B. Ronning, N. M. Pedersen, V. Bertelsen, E. Stang and I. H. Madhus: The oncoprotein ErbB3 is endocytosed in the absence of added ligand in a clathrin-dependent manner. *Carcinogenesis*, 33(5), 1031-9 (2012)
DOI: 10.1093/carcin/bgs128
PMid:22436610
52. L. Henriksen, M. V. Grandal, S. L. Knudsen, B. van Deurs and L. M. Grovdal: Internalization mechanisms of the epidermal growth factor receptor after activation with different ligands. *PLoS One*, 8(3), e58148 (2013)
DOI: 10.1371/journal.pone.0058148
PMid:23472148 PMCID:PMC3589378
53. K. Roepstorff, L. Grovdal, M. Grandal, M. Lerdrup and B. van Deurs: Endocytic downregulation of ErbB receptors: mechanisms and relevance in cancer. *Histochem Cell Biol*, 129(5), 563-78 (2008)
DOI: 10.1007/s00418-008-0401-3
PMid:18288481 PMCID:PMC2323030
54. M. Zhen, R. Heinlein, D. Jones, S. Jentsch and E. P. Candido: The ubc-2 gene of *Caenorhabditis elegans* encodes a ubiquitin-conjugating enzyme involved in selective protein degradation. *Mol Cell Biol*, 13(3), 1371-7 (1993)
DOI: 10.1128/MCB.13.3.1371
PMid:8441382 PMCID:PMC359446
55. M. R. Rogel, A. Jaitovich and K. M. Ridge: The role of the ubiquitin proteasome pathway in keratin intermediate filament protein degradation. *Proc Am Thorac Soc*, 7(1), 71-6 (2010)
DOI: 10.1513/pats.200908-089JS
PMid:20160151 PMCID:PMC3137152
56. A. G. Manford, C. J. Stefan, H. L. Yuan, J. A. Macgurn and S. D. Emr: ER-to-plasma membrane tethering proteins regulate cell signaling and ER morphology. *Dev Cell*, 23(6), 1129-40 (2012)
DOI: 10.1016/j.devcel.2012.11.004
PMid:23237950
57. F. Alber, S. Dokudovskaya, L. M. Veenhoff, W. Zhang, J. Kipper, D. Devos, A. Suprpto, O. Karni-Schmidt, R. Williams, B. T. Chait, A. Sali and M. P. Rout: The molecular architecture of the nuclear pore complex. *Nature*, 450(7170), 695-701 (2007)
DOI: 10.1038/nature06405
PMid:18046406
58. B. A. Hocevar, C. Prunier and P. H. Howe: Disabled-2 (Dab2) mediates transforming growth factor beta (TGFbeta)-stimulated fibronectin synthesis through TGFbeta-activated kinase 1 and activation of the JNK pathway. *J Biol Chem*, 280(27), 25920-7 (2005)
DOI: 10.1074/jbc.M501150200
PMid:15894542
59. J. Kang, Y. Shi, B. Xiang, B. Qu, W. Su, M. Zhu, M. Zhang, G. Bao, F. Wang, X. Zhang, R. Yang, F. Fan, X. Chen, G. Pei and L. Ma: A nuclear function of beta-arrestin1 in GPCR signaling: regulation of histone acetylation and gene transcription. *Cell*, 123(5), 833-47 (2005)
DOI: 10.1016/j.cell.2005.09.011
PMid:16325578
60. T. M. Brand, M. Iida, C. Li and D. L. Wheeler: The nuclear epidermal growth factor receptor signaling network and its role in cancer. *Discov Med*, 12(66), 419-32 (2011)
PMid:22127113 PMCID:PMC3305885

61. Y. N. Wang, H. H. Lee, H. J. Lee, Y. Du, H. Yamaguchi and M. C. Hung: Membrane-bound trafficking regulates nuclear transport of integral epidermal growth factor receptor (EGFR) and ErbB-2. *J Biol Chem*, 287(20), 16869-79 (2012)
DOI: 10.1074/jbc.M111.314799
PMid:22451678 PMCID:PMC3351284
62. D. M. Bryant, F. G. Wylie and J. L. Stow: Regulation of endocytosis, nuclear translocation, and signaling of fibroblast growth factor receptor 1 by E-cadherin. *Mol Biol Cell*, 16(1), 14-23 (2005)
DOI: 10.1091/mbc.E04-09-0845
PMid:15509650 PMCID:PMC539147
63. A. V. Vieira, C. Lamaze and S. L. Schmid: Control of EGF receptor signaling by clathrin-mediated endocytosis. *Science*, 274(5295), 2086-9 (1996)
DOI: 10.1126/science.274.5295.2086
PMid:8953040
64. H. M. Jopling, A. F. Odell, C. Pellet-Many, A. M. Latham, P. Frankel, A. Sivaprasadarao, J. H. Walker, I. C. Zachary and S. Ponnambalam: Endosome-to-Plasma Membrane Recycling of VEGFR2 Receptor Tyrosine Kinase Regulates Endothelial Function and Blood Vessel Formation. *Cells*, 3(2), 363-85 (2014)
DOI: 10.3390/cells3020363
PMid:24785348 PMCID:PMC4092869
65. I. Domingues, J. Rino, J. A. Demmers, P. de Lanerolle and S. C. Santos: VEGFR2 translocates to the nucleus to regulate its own transcription. *PLoS One*, 6(9), e25668 (2011)
DOI: 10.1371/journal.pone.0025668
PMid:21980525 PMCID:PMC3182252

activating protein; μ HD: μ -homology domain; VCA: Verprolin, cofilin, acidic; WASP: Wiskott-Aldrich syndrome protein; WAVE: WASP family verproline-homologous protein. WH1/2, WASP-Homology 1/2

Key Words: F-BAR proteins, Membrane receptor, Cellular trafficking, Nuclear translocation, Endocytosis, Review

Send correspondence to: Hong Wang, Centers for Metabolic Disease Research, Cardiovascular Research and Thrombosis Research, Temple University School of Medicine, MERB-1060, 3500 North Broad Street, Philadelphia, PA 19140, Tel: 215-707-5986, Fax: 215-707-5737, E-mail: hongw@temple.edu

Abbreviation: CIP4: CDC42-interacting protein 4; CRIB: CDC42-Rac interactive binding; CP, Cytoplasmic; CSK, cytoskeleton; ECM: Extracellular matrix; F-actin: Filamentous actin; F-BAR: Fes/CIP4 homology-Bin/Amphiphysin/Rvs; FCHO: FCH only; FCHSD: FCH and double SH3 domain proteins; FER: FES related; FX: F-BAR extension; GAS7: Growth arrest-specific 7; HR1: Protein kinase C-related kinase homology region 1; NOSTRIN: Nitric oxide synthase traffic inducer; N-WASP: Neural Wiskott-Aldrich syndrome protein; NC, Nucleus; PACSIN: Protein kinase C and casein kinase 2 substrates in neurons; PH, Pleckstrin homology; PM, plasma membrane; RhoGAP: Rho GTPase-activating protein; SH2: Src homology-2; SH3: Src homology-3; srGAP: Slit-Robo GTPase-