

## Environmental pollutants: an immunoendocrine perspective on phthalates

Margarita Isabel Palacios-Arreola<sup>1</sup>, Jorge Morales-Montor<sup>2</sup>, Cintia Jocelyn Cazares-Martinez<sup>3</sup>, Sandra Gomez-Arroyo<sup>3</sup>, Karen Elizabeth Nava-Castro<sup>3</sup>

<sup>1</sup>Laboratorio de Especiacion Quimica de Aerosoles Organicos Atmosfericos y Desarrollo de Tecnologias Verdes, Departamento de Ciencias Ambientales, Centro de Ciencias de la Atmosfera, Universidad Nacional Autonoma de Mexico, CP 04510, Ciudad de Mexico, Mexico, <sup>2</sup>Departamento de Inmunologia, Instituto de Investigaciones Biomedicas, Universidad Nacional Autonoma de Mexico, AP 70228, Ciudad de Mexico, CP, 04510, Mexico, <sup>3</sup>Laboratorio de Genotoxicologia y Mutagenesis Ambientales, Departamento de Ciencias Ambientales. Centro de Ciencias de la Atmosfera, Universidad Nacional Autonoma de Mexico, CP 04510, Ciudad de Mexico, Mexico

### TABLE OF CONTENTS

1. Abstract
2. Introduction
  - 2.1. Phthalates
    - 2.1.1. Phthalates as endocrine disrupting compounds (EDCs)
    - 2.1.2. Phthalate metabolism
  - 2.2. Molecular signaling induced by phthalates
    - 2.2.1. Estrogen receptor (ER) pathways
    - 2.2.2. Androgen receptor (AR) pathway
    - 2.2.3. The peroxisome proliferator-activated receptor gamma (PPAR-gamma) pathway
    - 2.2.4. The Aryl hydrocarbon receptor (AhR) pathway
  - 2.3. The role of endocrine receptor pathways on the immune system
3. Phthalates as immunomodulators
  - 3.1. In vitro studies
  - 3.2. In vivo studies
  - 3.3. Epidemiological associations
4. Conclusions
5. Acknowledgments
6. References

### 1. ABSTRACT

Phthalates are endocrine disrupting compounds (EDCs) used as plasticizers in a wide array of daily-use products, from flooring and automotive parts to medical devices and are even present in the children's toys. Since these compounds are not covalently bound other molecules, they leach from these synthetic products, causing a high level of human exposure to them. EDCs exert several endocrine effects, most typically, reduced biosynthesis of the male

hormone, testosterone and disturbances in estrogen, androgen, PPAR-gamma and AhR that control complex immunoendocrine regulatory networks. Besides impacting the developmental processes and long-term adverse effects, since cells of the immune system express endocrine receptors, and synthesize and respond to several hormones and other endocrine ligands, phthalates also cause dysregulation of immune system.

## 2. INTRODUCTION

Phthalates are considered endocrine disrupting compounds (EDCs), which are molecules that interfere with hormonal homeostasis. Particularly, phthalates have been shown to interact with several receptors involved in estrogenic and androgenic pathways (1). It is important to note that even though the main target of EDCs is the endocrine system, they may affect other tissues. One of such tissues are cells from the immune system, which are greatly susceptible to endocrine modulation. This relationship was first recognized with epidemiological studies that found a higher incidence of some autoimmune diseases -such as rheumatoid arthritis and systemic erythematous lupus- in the female population (2, 3). Later, it was demonstrated that sex steroids regulate multiple immunological functions like lymphocyte maturation, cellular migration, expression of MHC molecules, and cytokine production and that this regulation is mediated by hormonal receptors expressed by immune cell populations (4).

Though some immunological effects of EDCs have been revised previously (5), this review aims to summarize the current evidence of immune dysregulation caused specifically by phthalates; and linking them with the acknowledged molecular mechanisms under an immunoendocrine perspective.

### 2.1. Phthalates

Phthalates are alkyl diesters of phthalic acid and are used as plasticizers in PVC products, as solvents and fixatives in personal care products and as additives in the enteric coating of some drug tablets. Thus, phthalates are present in a wide array of daily-use products, from flooring and automotive parts to medical devices and even children's toys. Since these compounds are not covalently bound to the products, they easily migrate from them (6), leading to high human exposure.

Phthalate exposure has been estimated by biomonitoring studies and ranges between 1 and 2 ug/kg bw/day of a single compound for adults (7, 8) and up to 4 ug/kg bw/day for children (9). While oral

consumption of contaminated food and dermal absorption from personal care products are considered the main exposure routes (10–12), phthalates are ubiquitously found in environmental matrices such as in the atmosphere, soil and water bodies (13). Of particular interest is the presence of phthalates in the atmospheric aerosol, forming part of the organic content of fine particulate matter (PM<sub>2.5</sub>). Phthalate concentrations can range from 1 to 100 ng/m<sup>3</sup> outdoors (14, 15) and up to 1,000 ng/m<sup>3</sup> indoors (16, 17), being Bis-(2-ethylhexyl) phthalate (DEHP) and Di-n-Butyl phthalate (DBP) the most common.


#### 2.1.1. Phthalates as endocrine disrupting compounds (EDCs)

An Endocrine disruptor compounds is defined by the United States Environmental Protection Agency (EPA) as an “exogenous agent that interferes with synthesis, secretion, transport, metabolism, binding action, or elimination of natural blood-borne hormones that are present in the body and are responsible for homeostasis, reproduction, and developmental process” (18). EDCs mimic endogenous hormones and bind to their receptors inducing functional and regulatory activities of gene expression.

Phthalates exert sexually dimorphic effects, affecting in different ways to male and female individuals. Bases on epidemiological data, elevated levels of DEHP urinary metabolites were linked to female patients experiencing premature thelarche (19). On the other hand, phthalates effects on males are grouped into the so called “phthalate syndrome”, which is a type of testicular dysgenesis featuring cryptorchidism hypospadias, incomplete testicular descend, reduced ano-genital distance, low sperm count and/or quality, elevated risk of infertility and testicular cancer (6, 20–22), caused by a reduced testosterone synthesis.

#### 2.1.2. Phthalate metabolism

Phthalates are readily metabolized once in the organism, leading to the urinary and fecal excretion of their metabolites. The first metabolic phase is an enzymatic hydrolysis of one of the side chains, which turns the original diester into a monoester (e.g. DEHP into MEHP). Phase I is



**Figure 1.** Molecular signaling induced by phthalates. Phthalates and some phthalate metabolites may exert fast non-transcriptional effects by engaging and activating G protein-coupled receptors (GPCR), which activate downstream pathways that lead to intracellular calcium increases and activation of diverse protein kinases. Phthalates and their metabolites readily cross cell membranes, entering the cytoplasm. Once in the cytoplasm, they may bind to estrogen receptors (ER) and induce their dimerization and therefore their activation, leading to transcriptional activity, which may vary depending on the existing coregulators. Phthalates may also bind androgen receptor (AR), triggering transcriptional effects and/or regulating the activation of kinases and transcription factors. Aryl hydrocarbon receptor (AhR) may also bind phthalates, enter the nucleus and regulate transcription of responsive genes. Upon nucleus localization, phthalates may also act as Peroxisome proliferation-activated receptor gamma (PPAR $\gamma$ ) ligands, leading to transcriptional activity.

catalyzed by esterases, which are present in blood plasma and most tissues (23). It is important to note that phthalate monoesters can also exert biological effects and maintain binding potential to some receptors (24–27). After this first phase, the monoesters can be further transformed by hydroxylation and/or carboxylation of the remaining side chain, before being further metabolized to glucuronide conjugates in phase II (23).

## 2.2. Molecular signaling induced by phthalates

The biological action induced by phthalates differ from the target cell in which they have an effect. The best described mechanisms involve signaling through the hormone receptors: estrogen (ER) and androgen (AR) receptors; and by the peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ), although other non-transcriptional mechanisms have been described, such as the activation of G protein-coupled receptors (Figure 1).

### 2.2.1. Estrogen receptor (ER) pathways

Classical nuclear estrogen receptors (ER $\alpha$  and ER $\beta$ ) belong to the ligand-activated transcription factor family. These receptors are widely expressed in mammal tissues, some of them expressing ER $\alpha$  (uterus, liver, kidney and heart) or ER $\beta$  (ovary, prostate, hematopoietic and central nervous systems) and some others expressing both (mammary gland, thyroid and adrenal glands and some brain regions). Even though both receptors are highly homologous and display a similar affinity for their natural ligand 17 $\beta$ -estradiol and for estrogen response elements (EREs) in the DNA, (28, 29), ligand binding to one receptor or another triggers different responses. The key to such differential responses lays in the signaling pathways of ERs.

Estrogen receptors reside in the cytoplasm and form homo- or heterodimers upon ligand binding, translocate to the nucleus and bind to the EREs in the DNA, typically located near the promoter regions

## Phthalates as immunomodulators

of target genes. This activation leads to the recruitment of coregulators, which associate to the ER dimers, forming the initial transcriptional complex (30). It is important to note that coregulators can either increase a gene's transcription (coactivator) or inhibit it (corepressor) and that recruitment of one or another depends on the type of ER dimer (ER-alpha or ER-beta homodimers or heterodimer) and, of course, on which coregulators are present in a particular moment in a particular cell type.

The activity of estrogen receptors is not limited to target genes containing EREs. Ligand-activated ERs can modulate gene expression by attaching to other transcription factors which in turn act on their respective response elements (30). In addition to the former, estrogen receptors may also be embedded into the plasma membrane, where they can bind extracellular ligands. Moreover, some non-classical membrane estrogen receptors have been described, such as GPER (or Gpr30), which is a G protein-coupled receptor. These membrane-bound receptors trigger rapid effects such as calcium flux, second messenger production (cAMP, DAG, IP3) and kinase activation (31, 32).

Considering the former, estrogens effects depend not only on the dose, but on the homo- or heterodimers formation, their target (either nucleus or cytoplasmic proteins), the coregulators and active transcription factors existing and all of it varies according to the context and specific cell type.

### 2.2.2. Androgen receptor (AR) pathway

Similar to estrogen receptors, androgen receptor (AR) has a widespread distribution, from reproductive tissues to muscle, bone, adipose tissue, and cardiovascular, immune and nervous system (33). Endogenous ligands for AR are testosterone and dihydrotestosterone (DHT), a metabolite of the former with a greater agonistic activity.

AR resides in cytoplasm, attached to the cytoskeleton by associating with heat shock proteins or other chaperones. Upon androgen binding, AR suffers conformational changes that dissociate it from the chaperones, which allows it to interact with coregulators and translocate to the nucleus, where it can dimerize and bind to DNA regions denominated

androgen response elements (AREs). Once there, AR together with coregulators forms an initial transcription factor that recruits the whole transcription machinery (33, 34).

Androgens can also indirectly modulate genetic expression without binding to AREs. The ligand-AR complex can activate other signaling pathways such as ERK and MAPK or even sequester other transcription factors (e.g. AP-1), repressing in this way the expression of their target genes (33).

### 2.2.3. The peroxisome proliferator-activated receptor gamma (PPAR-gamma) pathway

Phthalates have been also described as peroxisome proliferators, inducing hepatocyte peroxisome and cellular proliferation, which in some studies in mice and rats, have an impact in the incidence of hepatocellular adenomas. This effect is dependent on the activation of the nuclear receptor peroxisome proliferator activated receptor gamma (PPAR-gamma). This receptor belongs to the nuclear receptor subfamily of ligand-inducible transcription factors, which were originally described as key transcriptional regulator in the metabolism of glucose and fatty acids, lipid storage and insulin sensitivity (35–37). In fact, it is one of the main targets in the treatment of metabolic syndrome, insulin resistance and type 2 diabetes (38), since it is highly expressed in white and brown adipose tissues.

Constitutively, PPAR-gamma forms a heterodimer with the retinoid X receptor (RXR) and they bind to specific DNA sequences termed PPAR response elements (PPRE). In basal conditions, PPAR-gamma is bound to corepressor molecules such as the silencing mediator of retinoid and thyroid hormone receptors (SMRT), the nuclear receptor corepressor (NCoR) and some histone deacetylases (HDACs). Upon ligand binding, there is a conformational change in the PPAR-gamma protein that allows the dissociation of its repressors and the recruitment of its coactivators. Among these are the CREB-binding protein (CBP), the histone acetyl transferase (p300) and the PPAR-binding protein 4 (PBP). During adipocyte differentiation, changes in the recruitment of coactivators allow the formation of different transcription complexes that modify the

## Phthalates as immunomodulators

responses to metabolic stimuli and ensure the activation of specific signaling pathways (39).

### 2.2.4. The Aryl hydrocarbon receptor (AhR) pathway

The AhR was initially described when analyzing the effects of the herbicide 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) induced in exposed chemical workers. This receptor belongs to the family of the bHLH Pas domain transcription factors that are involved in sensing environmental changes. The canonical signaling pathway is induced when its ligand reaches the cytoplasm and bind to the AhR –which in the absence of ligand, is attached to actin filaments by the interaction with the chaperon proteins HSP90, AIP and p23. The ligand-AhR/HSP90/p23 complex then translocate to the nucleus, where it is released from this complex and heterodimerizes with the ARNT protein. This dimer binds to genomic regions containing the dioxin response element (DRE) and induce the transcription of genes such as *CYP1A1*, *CYP1A2*, *CYP1B1* and the AhR repressor (*AhRR*). The expression and function of these proteins regulate negatively this signaling pathway by three mechanisms: 1) AhR proteosomal degradation – mediated by the covalently union of ubiquitin-; 2) Ligand degradation by *CYP1A1* and 3) AhR/ARNT complex disruption by the repressor AhRR. This repressor is a protein similar to AhR but differs in two characteristics: it lacks its ligand-binding domain and contains a transactivation domain in the C-terminal domain that recruit other co-repressors (40, 41).

In addition to this canonical pathway, the AhR/ARNT complex can recruit other proteins and regulate either transcription or chromatin remodeling. Among these proteins, the SW1/SNF remodeling complex and the steroid receptor coactivator-1 (SRC-1), CBP/p300, p160/bHLH-PAS, NCoA2/GRIP1/TIF2, pCIP (p300/CPB/CoIntegrator associated protein) and RIP140 or the ATP-dependent chromatin remodeling components such as BRG-1 (40, 41).

There are other signaling pathways related to AhR activation. For example, the ligand TCDD increase  $Ca^{2+}$  concentration by activating the Src tyrosine kinase and the Focal adhesion kinase (FAK). These two pathways converge in MAPK activation

and therefore, there is also an induced migration, adhesion and inflammation-related functions such as prostaglandin and arachidonic production.

### 2.3. The role of endocrine receptor pathways on the immune system

The immune and endocrine systems are part of the neuroimmunoendocrine network, a recently recognized multisystem circuitry that regulates homeostasis. Regarding the immune system, its cellular components not only express endocrine receptors, but synthesize and respond to several hormones and other endocrine ligands. Some molecular pathways typically associated to the endocrine system play a developmental and regulatory role on the immune system, such as the thymus involution through age, the sexual dimorphism in immune response and the tolerogenic character of immune response during pregnancy.

Estrogen receptors are expressed in most immune cell populations, from hematopoietic precursors to mature B and T lymphocytes, NK cells, monocytes and dendritic cells (42). However, expression patterns are different, with some cells expressing preferentially ER-alpha (e.g. CD4<sup>+</sup> T cells) and other expressing higher levels of ER-beta (e.g. B cells) (43, 44). In homeostatic conditions, ER-alpha acts on myeloid and lymphoid precursors, inducing developmental pathways (42).

The developmental importance of ER-alpha is depicted by the thymic and splenic hypoplasia observed in ER-alpha-deficient mice (45). ER has a profound effect on T lymphocytes which is sensitive to the dose or physiological levels of estradiol, the natural ligand. A good example is the shift from Th1 response towards Th2 and Treg expansion due to high levels of estrogen, as occurs during pregnancy. B cells are deeply modulated by estrogen, both ER-alpha and ER-beta pathways are involved in maturation, function and survival of this population (42), so that females tend to display a greater humoral response than males (46). Differentiation and maturation of dendritic cells, as well as macrophage effector functions are also modulated by estrogen receptor pathways (44).

## Phthalates as immunomodulators

AR is expressed in several cell populations from both myeloid and lymphoid origin and has been involved in sexual immunological dimorphism and immunosuppression (47). The majority of the studies regard androgens and androgen receptor as suppressive or regulatory. AR pathways downregulate T cell and B cell proliferation, as well as the inflammatory response (48). Androgens also modulate the functions of dendritic cells, downregulating expression levels of MHC, HLA and costimulatory molecules, which further modulates T cell response (47). One exception to the predominantly suppressive role of androgens is the neutrophil subpopulation, since androgens (via AR) promotes their differentiation and maturation, probably by enhancing Granulocyte colony-stimulating factor (G-CSF) signaling (48). Though histamine release by mast cells appears to be downregulated by androgens, testosterone has been shown to induce IL-33 expression, which drives the generation of innate lymphoid cells and basophils, which in turn produce Th2 cytokines and drive IgE antibody switch (47). A very interesting observation is that IgE is found to be higher in young males over females affected by allergic rhinitis (49).

PPAR-gamma is expressed in lower amounts in some immune cells such as macrophages, eosinophils, dendritic cells, T and B lymphocytes; however, it has a critical role in both differentiation and activation of these cells in diseases that involve chronic inflammation such as atherosclerosis, inflammatory bowel disease and rheumatoid arthritis (50).

Expression of PPAR-gamma is differentially expressed in different subtypes of macrophages. Resting bone-marrow-derived macrophages express low levels of PPAR-gamma mRNA, whereas activated peritoneal macrophages express high levels of PPAR-gamma (51). However, peritoneal macrophages incubated with 15-dPGJ<sub>2</sub> – a PPAR-gamma agonist- showed morphological features typical of resting cells, even in the presence of IFN-gamma. In addition, it also downregulates iNOS expression and inhibit IFN-gamma-dependent nitrite production. This suggests that might inhibit the expression of genes that become upregulated during macrophage differentiation and activation (51). In

contrast, other studies report that activation of PPAR-gamma primes primary human monocytes into M2 differentiation, but it is not able to modify marker expression of differentiated M1 or M2 macrophages (52).

Eosinophils, an immune cell population related to allergic responses and asthma, express PPAR-alpha, PPAR-gamma and PPAR-delta. In fact, the administration of PPAR-gamma agonists (rosiglitazone or pioglitazone) in a murine model of allergic asthma, induce a reduction in the number and activation state of eosinophils in the airways, determined by the expression of IL-4, IL-5 and ovalbumin-specific IgE, which suggest that PPAR-gamma may be protective factor in the pathogenesis of the asthma (53). This correlate with *in vitro* experiments in which the presence of high concentrations of PPAR-gamma agonists reduce the eotaxin-dependent eosinophil migration; while low concentrations of this compounds increase migration (54). Interestingly, this effect is mediated by changes in calcium influx, but not by the expression of CCR3 or phosphorylation of p38 or ERK (55).

As mentioned for eosinophils, PPAR-gamma expression in dendritic cells have been related to allergic responses. Particularly, PPAR-gamma deficient CD11b<sup>+</sup> dendritic cells fail to migrate to the lung draining lymph nodes and therefore a reduced capacity to polarize naïve CD4<sup>+</sup> T cells toward a Th2 phenotype (56, 57). Interestingly, Tuna and collaborators mention that the effect of PPAR-gamma activation depend on the tissue microenvironment. While in the lung it may induce a mucosal phenotype in mDCs and that loss of PPAR-gamma promotes an inflammatory phenotype (determined by the ability of BM-DC to polarize CD4 T cells toward iTregs and to induce CCR9 – a mucosal homing receptor- on T cells); deficiency of this receptor showed no change in the frequency or phenotype of mDC in the colon. This suggest that the intestinal microenvironment can maintain the mucosal DC phenotype of via PPAR-gamma-independent mechanisms (58).

T cells express PPAR-gamma 1, but not PPAR-gamma 2 and it is upregulated on activated T cells. Interestingly, its activation through agonists

## Phthalates as immunomodulators

(such as 15-deoxy- $\Delta$ 12, 14-prostaglandin J2 or troglitazone) inhibits PMA-mediated proliferation and induce a decrement in cell viability (59). This effect was also described in human and B cell lines. Incubation of B lineage cells with PPAR-gamma against but not with PPAR-alpha agonists, induce apoptosis as demonstrated by the increase of AnnexinV staining and the evaluation of DNA fragmentation in TUNEL assays (60).

Because of its function as an environmental sensor, AhR is widely expressed in the body and the immune system. However, it is highly expressed at the barrier sites such as the skin, lung and gut. In the immune system, the expression of AhR has been described in dendritic cells, B cells and some subpopulations of T cells such as Th17, and in lesser amount, in regulatory T cells (61).

In the T cell subpopulations, AhR is highly expressed in Th17 and in a lesser amount in regulatory T cells (Tregs), but it has not been found in naïve, Th1 or Th2 cells (41). The role of this receptor in Th17 cells remains mostly elusive, but some studies suggest that AhR interacts with NF- $\kappa$ B and STAT1, inhibiting those proinflammatory pathways (62) and promoting the Th17 phenotype by enabling cells to produce high levels of IL-17 and IL-22 (63). Regarding B cells, it has been reported that AhR is expressed in several subpopulations, although at different levels (41). AhR role in B cell function is still not well understood, but an *in vitro* study reported B cell maturation impairment and antibody production suppression when LPS stimulation occurred in presence of a potent AhR ligand (64). AhR is also expressed by antigen presenting cells, such as dendritic cells, whose differentiation and maturation can be affected by AhR activation. Furthermore, activation of AhR decreases the expression of class II MHC, as well as costimulatory molecules, promoting regulatory T cells differentiation (65).

## 3. PHTHALATES AS IMMUNOMODULATORS

### 3.1. *In vitro* studies

As summarized in Table 1, most studies about the impact of phthalates in the immune system are developed in murine macrophages either cell

lines or in primary cultures of peritoneal exudates. In these cells, it was found that some phthalates such as DEHP, DEP and MEHP promoted a pro-inflammatory state characterized by an increase in the expression and secretion of IL-1, IL-6, TNF-alpha, and the chemokine CXCL1 and ROS (66). Other studies found that the secretion of IL-6, IL-10 and the chemokine CXCL8 was enhanced, while TNF-alpha was impaired by DEP and DnBP presence during stimulation (67). They are also capable of increase the phagocytic index of rabbit lung macrophages, which is accompanied by an increment in the release of lysosomal hydrolases (68). However, Shertzer found that even when the phagocytic index after the infection with *S. aureus* was increased, there was a reduction in the rate of pathogen destruction; which may impact as a deficient response in subsequent exposures (69). On the contrary, other phthalates such as the DBP, BzBP, DEP and DPrP inhibit the production of IL-1beta, IL-6 and IFN-beta (70, 71), apparently by the inactivation of the IFN-beta promoter (70).

In T cells, phthalates seem to promote the polarization to the Th2 phenotype. Hansen *et al.* showed that 100  $\mu$ M of DEP and DnBP were capable to suppress the secretion of IL-2 and IL-4, TNF-alpha and IFN-gamma, with no impact in cell viability. It was found that DEHP stimulation might also increase the expression of TCR and CD3 molecules and antigen-induced proliferation (72). In total splenocyte cultures, DEHP significantly increased IL-4 and IFN-gamma concentrations, while MEHP increased IL-6 in supernatants. Interestingly, the effect of DEHP was mediated by the activation of the Ca<sup>2+</sup>/CaN/NF-AT signaling pathway, which was not observed with MEHP. These data suggest that both phthalates may induce the same outcome but by altering different cellular mechanisms (73). DEHP also induces B cell proliferation; activate Ca<sup>2+</sup> flux, membrane PKC translocation, ERK1/2 phosphorylation and NF-kappa B activation after 2 hrs (74).

In murine thymocytes, DEHP also increases B cell proliferation as demonstrated by Yamashita *et al.* (75). In contrast, MEHP was found to suppress B cell ((3)H) thymidine incorporation in both bone marrow-derived B cells and the B cell line BU-11 at doses of 25 or 100  $\mu$ M. This was explained

## Phthalates as immunomodulators

**Table 1.** *In vitro* studies in animal cells

Country	Cellular type	Immune system component	Phthalate	Findings	Reference
USA	Primary culture of rat testicle germ cell	Macrophages, IL-6, TNF-alpha and CXCL1	DEHP and DEP (100 uM up to 7 days)	The expression of proinflammatory cytokines (IL-6, TNF-alpha and CXCL1) are increased	(66)
USA	Rabbit alveolar macrophages	Macrophages	DEHP (2.5 uL per ml of serum)	Significant stimulation of phagocytic activity in macrophages pre-exposed to DEHP as well as release of lysosomal hydrolases.	(68)
USA	Rabbit lung macrophages	Macrophages	DEHP (2.5 uL per ml of serum)	The cultures previously exposed to DEHP, after infection with <i>S. Aureus</i> , show an increase in the phagocytic index and a decrease between 2 and 14 times of the rate of destruction of the pathogen.	(69)
Japan	RAW 264	IFN-beta	BBP, DBP, DOP, DEP, DHP, DPP and DPrP (50-200 uM)	Inhibition of IL-1beta and IL-6. Only DEP and DPrP inhibit the LPS-induced activation of IFN-beta promoter.	(70)
Japan	Mice (abdominal macrophages) and RAW 264 cell line	TNF-alpha	BBP (10 ng/mL)	Inhibits the production of TNF-alpha in RAW 264. In abdominal macrophages no response was seen.	(71)
Japan	Bone marrow-derived dendritic cells (BMDC) and splenocytes from mouse	Dendritic cells, splenocytes, IL-4 and IFN-gamma	DEHP (0.1-10 uM)	Increases BMDC differentiation but not the activation. Higher IL-4 and IFN-gamma expression on T lymphocytes in response to BMDC differentiation. Increased Th2 response in splenocytes.	(72)
China	Primary cultures of mouse splenocytes	IL-4, IL-6 and IFN-gamma	DEHP (1-50 uM) and MEHP (5-80 uM)	DEHP: Increases IL-4 and IFN- $\gamma$ , decreases the Th1/Th2 ratio (IFN-gamma / IL-4). MEHP: decreases the Th1/Th2 ratio (IFN-gamma / IL-6).	(73)
Korea	Primary cultures of mouse splenocytes	B lymphocytes	DEHP (100 uM)	Increases B cell proliferation by inducing Ca <sup>2+</sup> flux, PKC, ERK1/2 and NF-kappa B activation.	(74)
Japan	Mouse thymocytes and splenocytes	Thymocytes (IL-3, IL-4 and IFN-gamma) and splenocytes	DEHP (1 nM-1uM)	Stimulates the proliferative response in thymocytes and splenocytes. Increases the production of IL-3 in thymocytes, IL-4 and IFN-gamma in Con A-treated thymocytes	(75)
USA	Mouse bone marrow-derived B cells and BU-11 cell line	B lymphocytes	MEHP (10-200 uM)	Induces apoptosis and suppresses B cell proliferation in developing bone marrow	(76)
USA	PD31 cell line (murine pre-B cells) and primary cultures of rainbow trout kidney	B cells, Plasmatic cells and IgM	DEHP (1-256 uM)	DEHP: Inhibits B cells proliferation and reduces the number of plasmatic cells secretory of IgM. Accelerates the terminal differentiation of B cells	(77)
Japan	Rat basophils RBL-2H3	Mast cells	DBP, DIBP and DEHP (50-500 uM)	Without Ag stimulation, the phthalates don't cause a significant increase in degranulation. In cultures sensitized with Anti IgE, the three induce degranulation: DIBP > DBP > DEHP.	(78)
Korea	RAW 264.7 cell line (murine macrophages)	iNOS, TNF- $\alpha$ and IL-1beta	DCHP, DEP and DBP (2-200 uM)	DCHP, DEP, DBP: changes in iNOS mRNA expression. DBP: Suppression of TNF-alpha and IL-1beta transcription	(80)

contd...

## Phthalates as immunomodulators

**Table 1.** Contd...

Country	Cellular type	Immune system component	Phthalate	Findings	Reference
Korea	RAW 264.7	Macrophages, iNOS	BBP (2-200 uM)	Reduces the production of NO via regulation of iNOS mRNA expression	(81)
China	Macrophages of murine peritoneal exudate	IL-1 $\beta$ , IL-6, IL-10, IL-12 and TNF-alpha	DBP (1-100 uM)	Inhibition of IL-1beta, IL-6 and TNF-alpha production. No change in IL-10 and IL-12 synthesis	(82)
Norway	(RAW 264.7)	TNF-alpha	MEHP (0.3-1 mM)	Increases TNF-alpha and changes the cellular morphology of macrophages to a more elongated form, reminiscent of differentiated M2 macrophages	(83)
Japan	Peripheral blood mononuclear cells (PBMC) of mice	Mature and immature DCs	DEHP and MEHP (0.1-10 uM)	DEHP inhibits the expression of DC differentiation and maturation markers. MEHP has no effect.	(84)
Norway	Primary cultures of rat alveolar macrophages	TNF-alpha, ROS, CXCL2 and leukotriene B4 (LTB <sub>4</sub> )	MEHP (0.1 – 1 mM)	MEHP increases the production TNF-alpha, ROS, CXCL2 and (LTB <sub>4</sub> )	(85)
Finland	RAW 264.7	ROS production and cell death	DEHP, DBP, BBP and DIBP (1 nM – 1 mM)	Increased ROS production. Only DBP and DBIP caused cell death due to necrosis at their highest doses (effect>with DIBP).	(86)
USA	primary cultures of murine cells	IL-4, IFN-gamma and IgE	DEHP and DINP (0.1–10 uM)	By themselves they don't induce the production of IL-4 and IFN-gamma in mouse lymph nodes. In stimulated cells with PMA or metalloproteins an increase in IL-4 is observed.	(87)
UK	Human and rat neutrophils	CD11b (lymphocyte activation marker)	DEHP (1-3 mg/L)	DEHP initiates inflammatory response in human and rat neutrophils (by expression of CD11b)	(88)
Japan	Carp leukocytes	Leukocytes and kidney macrophages	DEHP and DBP (1–1000nM)	Both regulate the function of carp phagocytic cells	(89)
UK	Rat peritoneal leukocytes	Leukocytes	DEHP, MEHP and DBP (1–100 ug/mL)	Inhibition of the formation of prostaglandins and leukotrienes (PGE-2 y LTB <sub>4</sub> ) in leukocytes.	(90)

by an increase in B-cell apoptosis at high doses and an apparent arrest without death at low doses (76).

Interestingly, these alterations are also seen in other organisms such as the rainbow trout, in which, *in vitro* DEHP exposition of B cells inhibited in a dose-dependent way the proliferation of these cells. Also, authors found that there were changes in the differentiation that generates deficient plasmablast expansion and a reduced number of IgM-secreting plasma cells (77).

Regarding to other immune cell types, there is also evidence in which the effect of some phthalates modifies the function of bone marrow-derived dendritic

cells (BM-DC) and basophils. In DC's, the presence of 10 uM of DEHP during the differentiation of BM-derived macrophages increased the expression of maturation markers such as MHC-II, CD80/86, CD11c and DEC205. It also increased their capacity to induce secretion of IFN-gamma, IL-4 and IL-10 by T cells. This effect was also observed when DC's were differentiated in normal conditions and DEHP was added during the activation. Interestingly, at 100 uM of DEHP during activation, the percentage of maturation/activations markers were reduced and their antigen presenting activity is not affected as observed by the reduced proliferation of co-cultured T cells. Authors suggest that the enhancement on these markers and function might contribute to the aggravating effect of DEHP on allergic

## Phthalates as immunomodulators

disorders, which also may be useful for the treatment in this kind of pathologies (72).

Finally, since air pollutants have been related to airway diseases, the study of immune cell populations related to allergies has gained relevance. The RBL-2H3 cells, a basophil-derived cell line, which has been described also as a mast cell model, was exposed to different phthalates - DPB, DiPB and DEHP- at concentrations of 50-500  $\mu$ M. Authors described that in the absence of antigen stimulation, none of those compounds induce  $\beta$ -hexosaminidase release. However, when cells were pre-sensitized and then stimulated with an antigen, all of the phthalates increased the degranulation of these cells (78). Another report found an interest correlation with the number of carbon atoms present in the phthalates and the histamine-release of human basophils. In this work, PMBC's were incubated in the presence different phthalates and the stimulated with any of the following stimuli: anti-IgE, the bacterial derived peptide fMLP, a calcium ionophore or an allergen (cat hair extract). As described before, there was no effect on the histamine release in the absence of activation. Interestingly, only the phthalates and monophthalates with an 8-carbon atoms alkyl chain length were the stronger histamine-release potentiators, when there were no differences for the 4-, 9- or 10-carbon chain (79).

Some studies have also assessed the effect of phthalates on human cell lines and primary cultures (Table 2). As in the former studies, heterogeneity in exposure times and concentrations is observed, but some general features can be distinguished. It is interesting to note that, in general terms, a more reactive or proinflammatory response is observed in human cells *in vitro* exposed to different phthalates, regardless of the heterogeneity of models and exposure schemes.

Studies with THP, a macrophage cell line seem to concur in that phthalates and their metabolites increase the cytokine and chemokine production (79, 91, 92). Tetz and cols. Also observed a proinflammatory effect of phthalate exposure, with activation of COX pathway and increased prostaglandin production (93). An interesting study

by Teixeira noted that the effect of phthalate exposure showed differential features, depending on the macrophage phenotype, either M1 or M2 (94).

One of the first studies that documented the effect of phthalates on immune cells was derived from the concern of donated blood being storage in plastic bags; this study found decreased chemotaxis and bactericidal capacity of neutrophils (95). However, more recent studies using human granulocytes have found rather an increase in chemotaxis, IL-8 (96, 97), and promotion of inflammatory response (88).

Mast cells are of particular interest, given their role in allergic disease. Lee and cols. Reported an increase in IFN-gamma and IL-4 production upon DEHP exposure, which is linked to a more reactive phenotype. However, more studies regarding mast cells differentiation, migration and function are desirable.

### 3.2. *In vivo* studies

Given the complex multidirectional regulation that occurs between immune and endocrine system, it is clear that, although they offer valuable information, *in vitro* studies don't fully represent what occurs in a whole organism, and it is why *in vivo* studies are of such importance. Data from the full literature review is detailed in Table 3.

In general terms, phthalate exposure leads to higher immunoglobulin levels. In some models, an increase of IgG -mainly IgG, has been observed (106–110), while in sensitized animals is the IgE class which is augmented (111). For one of the most studied phthalates – DEHP, the increase in IgE seems to be mediated by the promotion of differentiation of B cells into plasma cells (112) and by the stimulation of follicular T helper cells (Thf), producing higher IL-21 and IL-4 (106).

In diverse pathologies that are characterized by hypersensitivity, such as asthma, dermatitis and rhinitis, phthalate exposure has been shown to increase Th2 (DBP, DPP, DEHP and DINP) and Th17 cytokines (DBP) (110, 113, 114). In animal asthma models, the most studied phthalates -DEHP

## Phthalates as immunomodulators

**Table 2.** *In vitro* studies with human cells

Country	Cellular type	Immune system component	Phthalate	Findings	Reference
Denmark	Monocytes/Macrophages and T cell (peripheral blood mononuclear cells)	Monocytes/Macrophages	DEP, DnBP and MnBP (Multiple concentrations)	DEP and DnBP increase IL-6, IL-10 and CXCL8. They decrease the synthesis of TNF- $\alpha$ . <b>T cells:</b> decrease in IL-2, IL-4, TNF-alpha and IFN-gamma expression.	(67)
Denmark	THP-1 and peripheral blood mononuclear cells (PBMC) of allergic and healthy individuals	THP-1: IL-1 $\beta$ , IL-6, IL-12 $\alpha$ PBMC: IL-4, IL-5 and IFN-alpha	MBUP, MBEP, MEHP, MOP, MINP and MIDP. (0.2-200 ug/mL)	Mono-phthalates don't increase the cytokine production in TPH-1 alone but increase it upon LPS stimulation. There seems to be a relationship between the length of the side chains and the toxicity. MBUP tends to increase the levels of IL-4 in PBMC of allergic patients	(79)
UK	Human and rat neutrophils	CD11b (lymphocyte activation marker)	DEHP (1-3 mg/mL)	DEHP initiates inflammatory response in neutrophils (by expression of CD11b) of humans and rats	(88)
Slovak Republic	THP-1	TNF-alpha, IL-1beta, IL-8	DiP (0.2-10 uM)	At high doses (10 uM), increases IL-8 secretion	(91)
France	THP-1 cell line (differentiated macrophages)	IL-1 $\beta$ , TNF- $\alpha$ and IL-8 (secretion of cytokines induced by LPS)	DEHP and DBP (0.001-10 uM)	DBP: increased expression of TNF-alpha	(92)
USA	THP-1 and placental macrophages (PM)	COX-2 and prostaglandin E2	MEHP (10-180 uM)	MEHP: Increases the release of PGE2 via increased expression of COX2	(93)
Portugal	Peripheral blood-derived M1 and M2 macrophages	IL-1beta, IL-6 and IL-10	DEHP and DBP (1 uM)	M1 macrophages: DEHP—>Increase the expression of IL-1beta, IL-10 and decrease IL-6. DBP—> Increase the expression of IL-1 $\beta$ is not affected. IL-10 and IL-6 the same DEHP. M2 macrophages: DEHP—>IL-1beta is not affected. IL-10 and IL-6 decrease. DBP—>Increase the expression of IL-1beta. IL-10 and IL-6 decrease.	(94)
Japan	Peripheral blood leukocytes	Granulocytes	DEHP (43.4 ug/cm <sup>2</sup> in blood storage bags)	Dose-dependent reduction of chemotaxis and bactericidal. Cell count and phagocytic activity aren't modified.	(95)
Italy	Primary cultures of granulocytes isolated from blood	Chemotherapy and ROS production	DEHP (0.0002-0.0024 %)	Stimulates chemotaxis and ROS production	(96)
Puerto Rico	TK6 cell line (Lymphoblasts)	Mitochondrial membrane permeability, ROS, Caspases 3 y 7	DEHP and MEHP (10-500 uM)	Both compounds affect the viability of lymphoblasts by increasing the permeability of the mitochondrial membrane, generating ROS and activating Caspases 3 and 7.	(97)
Norway	Peripheral blood mononuclear leukocytes	Leukocytes	TBEP and DEHP (0.1 uM-0.1M)	Affects the binding of beta-adrenergic ligand in mononuclear leukocytes (interacts with specific and nonspecific binding sites).	(98)
Germany	Peripheral blood leukocytes	Leukocytes	DBP and DiBP (354 mM)	Genotoxicity: DiBP >DEHP	(99)
Germany	Peripheral blood lymphocytes	Lymphocytes	MEHP (0.1-2.5 mM)	Genotoxicity	(100)

contd...

## Phthalates as immunomodulators

**Table 2.** Contd...

Country	Cellular type	Immune system component	Phthalate	Findings	Reference
USA	Primary cultures of umbilical cord blood	IL-1beta, IL-8, IL-6 and VEGF	MEHP (100–500 uM)	Inhibits neutrophil apoptosis (neonatal cells are more sensitive than adults). In adults and neonates increases the production of VEGF and IL-1beta, only increases IL-8 in adult neutrophils. IL-6 is not affected.	(101)
Korea	HMC-1 cell line (mast cells)	TNF-alpha and IL-4	DEHP (100 uM)	Increases the expression of TNF-alpha and IL-4	(102)
Japan	THP-1	TNF-alpha, IL-1beta, IL-8 and IL-6	DEHP (200uM)	Increases TNF-alpha, IL-1b, IL-8 and IL-6 secretion. Increases the mRNA expression levels of IL-8, CXCL1, CXCL2, CXCL3, CXCL6, CCL3, MMP10, CSF2, TNF-a, IL-1beta and IL-6. Induces p65 NF-kappa B translocation to the nucleus.	(103)
Taiwan	pDCs (Plasmacytoid dendritic cells), serum	IFN-alpha, IFN-beta, IFN-gamma and IL-13	DEHP and BBP (1–100 nM)	DEHP and BBP: suppress the expression of IFN-alpha, and IFN-beta. BBP: Suppresses the expression of IFN-gamma but increases the IL-13 (for T cell CD4+)	(104)
Taiwan	MDA-MB-231 cell line (adenocarcinoma)	mdDCs and TADCs (CXCL1)	BBP (1 and 10 uM)	BBP stimulates the production of CXCL1 by the TADCs	(105)

and DINP- have been shown to increase pulmonary inflammation, antibody production (specially IgE), and IL-4 production. This Th2 polarization has also been observed in an allergic rhinitis model, where DEHP exposure leads to higher IL-13 levels in the nasal epithelium (114). In cutaneous hypersensitivity models, DBP, DPP and DINP have been shown to promote Th2 response, as denoted by a high IL-4 production (115, 116).

Moreover, phthalate exposure also stimulates the production of chemoattractant molecules, such as TSLP, CCL17 and CCL22 (115, 117, 118), which coincides with another common finding: a higher recruitment of antigen presenting cells like dendritic cells and macrophages. This phenomenon, known as the adjuvant effect, has been observed in skin and other tissues, including adipose tissue, and for various phthalates (119, 120).

One of the most typical target organs regarding phthalate toxicity is the male gonad. The effect of phthalates on this tissue has been mainly attributed to pure endocrine mechanisms, but it seems that immunomodulatory mechanisms may as well play an important role. It has been observed that exposure

to some phthalates, fundamentally DEHP, causes an increased infiltration of macrophages and other leukocytes in the testis, together with a higher expression of proinflammatory cytokines, such as IL-1 $\beta$  and IFN-gamma, which leads to germinal cell apoptosis and lower testosterone levels (121–124).

### 3.3. Epidemiological associations

Epidemiological studies, albeit heterogeneous, support the immunomodulatory effects of phthalates, particularly due to prenatal exposure. Data from the full literature review is detailed in Table 4. A 70% higher risk of asthma diagnosis at age 5-11 was observed in children born to women with higher urinary levels of BBzP and DnBP metabolites during their pregnancies (132). Another study, carried out in a Spaniard cohort, also found an association between asthma diagnosis at age 7 and maternal urinary levels of MBzP and the sum of DEHP metabolites (133). In a similar fashion, a study in Taiwan reported an association between high maternal DEHP exposition and asthma incidence, particularly in boys (134). In a French cohort, a study found a correlation between maternal urinary concentrations of DiBP and DiNP metabolites and eczema occurrence; interestingly, this association was stronger for boys (135).

## Phthalates as immunomodulators

**Table 3.** *In vivo* studies in animal models

Country	Organism	Immune system component	Phthalate	Findings	Reference
China	Mice	IgE, IgG1, Tfh (follicular T helper), Plasmatic cells, IL-21 and IL-4	DEHP (30-300 ug/Kg, oral)	DEHP acts on Thf cells by increasing its production of IL-21 and IL-4, which leads an increase of IgE and IgG1 secretion.	(106)
Denmark	WT and PPAR- $\alpha$ -deficient mice	IgG1, IgE and IgG2a	DEHP (100 ug together with OVA, i.p.)	It induces increased levels of IgG and IgG2a in both strains of mice	(107)
Denmark	Mice	IgG1, IgE and IgG2a	DEHP (0.022-13 mg/m <sup>3</sup> , aerosol, with OVA, 5 days/week, 2 weeks)	DEHP is capable of increasing serum IgG1 levels and inflammatory cell levels in the lungs but only at very high concentrations	(108)
Denmark	Mice	IgG1, IgE, IgG2a, eosinophils and lymphocytes (serum, bronchoalveolar fluid and draining lymph nodes)	MEHP (0.03-0.4 mg/m <sup>3</sup> , aerosol, with OVA, 5 days/week, 2 weeks)	Increase in IgG1. > number of eosinophils and lymphocytes in bronchoalveolar fluid	(109)
Korea	Mouse	Cytokines, immunoglobulines, lung inflammation	DINP (50 mg/Kg, i.p. on days 0,3 and 10)	DINP suppresses Th1 polarization and promotes Th2. In asthmatic mice, DINP increases IL-4, IL-5, IgE, IgG1, decreases IgG2a and IFN-gamma (in serum) and induces lung inflammation s (Caspases 1 and 3 increased)	(110)
Sweden	Mice	IgE, IL-4, IFN-gamma	DEHP (30-3000 ug/kg/day, oral, 52 days.	In sensitized mice, serum IgE increases in a dose-dependent manner. IL-4 in lung tissue increases in sensitized mice exposed to the highest DEHP dose	(111)
USA	Salmon	B lymphocytes	DEHP, DMP, DBP, BBP, DEP, and DnOP (from plastic-polluted lake)	DEHP accelerates the differentiation towards the plasmatic cells (secretory of Ab)	(112)
China	Mice	IgE, IFN-gamma (Th1), IL-4 (Th2), IL-17A (Th17), TNF-alpha, IL-5 and IL-13	DBP (0.4-40 mg/Kg/day, 40 days)	In lesions due to atopic dermatitis, the exposure to DBP increases Th2 and Th17 cytokines	(113)
Japan	Mice with induced allergic rhinitis	IL-5, IL-6, IL-12 and IL-13	DEHP (0.0004-0.16 ug, nasal instillation)	Increased levels of nasal IL-13 in mice treated with the allergen +DEHP	(114)
Japan	Mice	CCL17, CCL22, IL-4	DINP (0.15-150 mg/Kg/day, i.p., days -5, 2, 9 and 16)	DINP increases the production of CCL17, CCL22 and IL-4 in induced atopic dermatitis	(115)

contd...

## Phthalates as immunomodulators

**Table 3.** Contd...

Country	Organism	Immune system component	Phthalate	Findings	Reference
Japan	Mice	IL-4	DBP and DPP (80 ul, together with acetone, as dermal vehicle)	Increased FITC <sup>+</sup> DC in lymph nodes. Increased IL-4 in draining lymph nodes after skin sensitization.	(116)
Japan	Mice with induced contact sensitivity	TSLP (Th2 response cytokine) and IL-4	DBP (Mixed 1:1 with acetone, used as vehicle, dermally administered)	Increases the expression of TSLP in sensitized skin. DBP is essential for sensitization with FITC. DBP induces migration of Ag presenting cells	(117)
USA	Mice	TSLP (cytokine related to IL-7) produced by epithelial cells	DBP (Mixed 1:1 with acetone, used as vehicle, dermally administered)	The epicutaneous application of DBP induced the expression of TSLP in the skin	(118)
Japan	Mice	Macrophages and DC	DBP and DPP (Mixed 1:1 with acetone, used as vehicle, dermally administered)	In the presence of these phthalates, a number of antigen-presenting DCs are observed in the lymph nodes (they migrate from the skin, presenting FITC)	(119)
Taiwan	Mice	Macrophages, TNF-alpha, IL-1 $\beta$ and IL-6	DEHP (500 mg/Kg/day, 4 weeks)	Promotes inflammation by increasing serum of TNF-alpha, IL-1 $\beta$ and IL-6. It also promotes the infiltration of macrophages in white adipose tissue.	(120)
USA	Rats	Macrophages	MEHP (1g/Kg, oral)	Induces the infiltration of CD11b+ macrophages in testis of immature peripubertal rats, triggering germ cell apoptosis. No infiltration observed in adult rats.	(121)
Japan	Mice	Lymphocytes, macrophages, IL-10 and IFN-gamma	DEHP (0.01-0.1 % in food)	Greater presence of lymphocytes and macrophages in testis, along with increased expression of IL-10 and IFN-gamma mRNA.	(122)
China	Rats	IL-1beta (of testicular macrophages)	DBP (50, 250 mg/Kg/day, 90 days)	Increases the expression of IL-1beta mRNA (and suppresses the production of testosterone)	(123)
Japan	Mice	IFN-gamma	DEHP (0.01% in food)	Increases the expression of IFN-gamma and the infiltration of leukocytes in autoimmune testicular inflammation.	(124)
New Zealand	Mice	Dendritic cells	DBP (Mixed 1:1 with acetone, used as vehicle, dermally administered)	Stimulation of Th2 response	(125)
China	Zebrafish	Macrophages	DBP (0.02 – 2.0 uM, embryo culture)	Decrease in macrophage formation.	(126)

contd...

## Phthalates as immunomodulators

**Table 3.** Contd...

Country	Organism	Immune system component	Phthalate	Findings	Reference
Japan	Mice with induced peritonitis	Macrophages, Neutrophils and IL-5	DEHP 1, 100 ug/mouse, i.p., 2/week, 4 times)	Exaggerates inflammation by increasing macrophages and neutrophils and increases IL-5 and eotaxin production.	(127)
Netherlands	Rats	TNF-alpha	DEHP (1, 1000 mg/Kg/day, oral, PND 10-50 or 50-90)	The exposure to DEHP increases TNF-alpha levels in splenocytes of adult rats (stimulated with LPS)	(128)
Japan	Epicutaneously sensitized mice	IL-4, IFN-gamma (from draining lymph nodes)	DEHP and DBP (Mixed 1:1 with acetone, used as vehicle, dermally administered)	DBP: Reduction of IL-4. DEHP: Reduction of IL-4 and increase of IFN-gamma	(129)
Japan	OVA-immunized mice	Splenocytes and thymocytes	DEHP (10 uM, drinking water, 4 weeks)	It has no effect on the production of Ab. Stimulates the proliferation of splenocytes treated with Con A (mitogen for T cells) and of the thymus cells treated with Con A.	(130)
USA	Rats	Kupffer cells	DEHP (1.2 g/Kg, oral)	An increase in free radicals in the bile of rats is observed after intragastric administration of DEHP. The increase in free radicals is due to the activation of Kupffer cells.	(131)

At the molecular level, Ashley-Martin didn't find an association between maternal MCPP (a nonspecific secondary metabolite) and levels of IgE, nor IL-33 in neonates (136). However, another study where MEP, MEHP, MBP and MBzP were measured, found a positive association between maternal urinary levels of those metabolites and IgE levels of children at age 2 and 5 (137). Interestingly, this association was only significant in boys.

Regarding current exposure in adults, the NHANES 2005-2006 study found an association between urinary levels of high molecular weight phthalate metabolites and allergic sensitization (defined by the response to at least 19 specific IgE antigens) (138). Furthermore, MBzP concentration correlated with the presence of allergic symptoms at the moment of the sampling. However, this same study found that in children there was an inverse relationship between urinary concentrations of phthalate metabolites and the incidence of asthma and hay fever. In another study, in adults allergic to dust mite, inhalation of DEHP-containing dust modified the expression of G-CSF, IL-5

and IL-6 in a differential fashion, depending on the concentration (139). At low levels, those cytokines increased, while at high concentrations both G-CSF and IL-6 decreased. It is interesting to note that, similarly to what is observed with some hormones, the dose-response relationship seems to be non-monotonic.

## 4. CONCLUSIONS

The immune system is regulated by a complex crosstalk between its own soluble mediators and several endocrine pathways. Immune cells not only express endocrine receptors but synthesize and respond to several hormones and other endocrine ligands. In this context, perturbations in the endocrine homeostasis may lead to immune dysregulation. Regarded as EDCs, phthalates and their metabolites interact with estrogen, androgen, PPAR-gamma and AhR pathways, and there is vast *in vitro* evidence of the effects these compounds exert on immune cells. Furthermore, reports on animal models confirm these phenomena occur in a whole organism setting. However, epidemiological evidence is the one that raises the more concerns, since it points out two

## Phthalates as immunomodulators

**Table 4.** Epidemiological studies on phthalate exposure

Country	Biological sample	Study cohort	Immune system component	Phthalate	Findings	Reference
USA	Urine of pregnant inner-city women	CCCEH (1998-2006)	Asthma diagnosis in children between 5 and 11 years old	BBzP, DnBP, DEHP, DEP	Risk of asthma was > 70% higher among children with maternal prenatal BBzP and DnBP metabolite concentrations in the third versus the first tertile.	(132)
Spain	Urine of pregnant women	INMA (2004-2008)	Atopy and asthma in progeny	Phthalate metabolites	Sum of DEHP metabolites, as well as MBzP during pregnancy increase the risk of asthma at age 7	(133)
Taiwan	Urine of pregnant women and children	Taiwan Maternal and Infant Cohort Study (2000-2009)	Asthma occurrence in progeny	Phthalate metabolites	Prenatal and postnatal exposure to phthalate was associated with the occurrence of asthma in children, particularly for boys	(134)
France	Urine of pregnant women	EDEN (2003-2006)	Development of Eczema in progeny	DiBP and DiNP metabolites	Occurrence of eczema in childhood is probably influenced by prenatal exposure to phthalates, in boys	(135)
Canada	Umbilical cord serum	MIREC (2008-2011)	IgE and IL-33	11 phthalate metabolites	Inverse non-linear associations between maternal urinary MCPP levels and elevated of IgE and IL-33 levels	(136)
Taiwan	Serum for children at 2 and 5 years	<i>Ad hoc</i> : from medical centers, regional hospitals, local hospitals, and clinics in Taiwan in 2004	IgE	MEP, MEHP, MBP and MBzP	Phthalate levels positively correlated with serum IgE levels only in boys.	(137)
USA	Urine	NHANES (2005-2006)	Allergic symptoms and sensitization	Phthalate metabolites	MBzP positively associated with allergic symptoms in adults. Mono-(3-carboxypropyl) phthalate and the sum of DEHP metabolites positively associated with allergic sensitization in adults. Phthalate metabolites inversely associated with asthma and hay fever in children.	(138)
Austria	Serum of healthy and allergic to home dust individuals	16 healthy and 16 HDM-allergic subjects	G-CSF, IL-5 and IL-6	DEHP	On allergic subjects, low concentrations of DEHP in the dust increased G-CSF, IL-5 and IL-6, while high concentrations of DEHP decreased G-CSF and IL-6.	(139)

relevant aspects of immune dysregulation associated to phthalates: developmental and long-term effects.

Taken together, the evidence suggests more consciousness should arise regarding the use of phthalates, especially in daily use personal care items intended for its use by vulnerable populations. We hope the recognition of the

immunoendocrine modulation network and the role it plays from early developmental stages will lead a more integrative perspective in toxicological studies. Regarding phthalates, more studies on immunological imprinting and potential neurological effects are desirable, since this system is also profoundly modulated by endocrine signaling.

## 5. ACKNOWLEDGMENTS

This work was supported by Grant No. IA202919 from Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica (PAPIIT) and Grant No. FC-2016-2125 from Consejo Nacional de Ciencia y Tecnología. Margarita Isabel Palacios-Arreola is a postdoctoral fellow from Dirección General de Asuntos del Personal Académico, UNAM.

## 6. REFERENCES

1. A Engel; T Buhrke; F Imber; S Jessel; A Seidel; W Völkel; A Lampen. Agonistic and antagonistic effects of phthalates and their urinary metabolites on the steroid hormone receptors Eralpha, Erbeta, and AR. *Toxicol Lett* 277, 54-63 (2017)  
DOI: 10.1016/j.toxlet.2017.05.028
2. GS Cooper; BC Stroehla. The epidemiology of autoimmune diseases. *Autoimmun Rev* 2, 119-125 (2003)  
DOI: 10.1016/S1568-9972(03)00006-5
3. A Bouman; MJ Heineman; MM Faas. Sex hormones and the immune response in humans. *Hum Reprod Update* 11, 411-423 (2005)  
DOI: 10.1093/humupd/dmi008
4. S Muñoz-Cruz; C Togno-Pierce; J Morales-Montor. Non-Reproductive Effects of Sex Steroids: Their Immunoregulatory Role. *Curr Top Med Chem* 11, 1714-1727 (2011)  
DOI: 10.2174/156802611796117630
5. K Nowak; E Jabłońska; W Ratajczak-Wrona. Immunomodulatory effects of synthetic endocrine disrupting chemicals on the development and functions of human immune cells. *Environ Int* 125, 350-364 (2019)  
DOI: 10.1016/j.envint.2019.01.078
6. SM Duty; AM Calafat; MJ Silva; L Ryan; R Hauser. Phthalate exposure and reproductive hormones in adult men. *Hum Reprod* 20, 604-610 (2005)  
DOI: 10.1093/humrep/deh656
7. M Wittassek; GA Wiesmüller; HM Koch; R Eckard; L Dobler; J Müller; J Angerer; C Schlüter. Internal phthalate exposure over the last two decades - A retrospective human biomonitoring study. *Int J Hyg Environ Health* 210, 319-333 (2007)  
DOI: 10.1016/j.ijheh.2007.01.037
8. HM Koch; H Drexler; J Angerer. An estimation of the daily intake of di(2-ethylhexyl) phthalate (DEHP) and other phthalates in the general population. *Int J Hyg Environ Health* 206, 77-83 (2003)  
DOI: 10.1078/1438-4639-00205
9. HM Koch; M Wittassek; T Brüning; J Angerer; U Heudorf. Exposure to phthalates in 5-6 years old primary school starters in Germany-A human biomonitoring study and a cumulative risk assessment. *Int J Hyg Environ Health* 214, 188-195 (2011)  
DOI: 10.1016/j.ijheh.2011.01.009
10. JD Meeker; S Sathyanarayana; SH Swan. Phthalates and other additives in plastics: human exposure and associated health outcomes. *Philos Trans R Soc B Biol Sci* 364, 2097-2113 (2009)  
DOI: 10.1098/rstb.2008.0268
11. JR Varshavsky; R Morello-Frosch; TJ Woodruff; AR Zota. Dietary sources of cumulative phthalates exposure among the U.S. general population in NHANES 2005-2014. *Environ Int* 115, 417-429 (2018)  
DOI: 10.1016/j.envint.2018.02.029

## Phthalates as immunomodulators

12. M Wittassek; HM Koch; J Angerer; T Brüning. Assessing exposure to phthalates - The human biomonitoring approach. *Mol Nutr Food Res* 55, 7-31 (2011)  
DOI: 10.1002/mnfr.201000121
13. A Bergé; M Cladière; J Gasperi; A Coursimault; B Tassin; R Moilleron. Meta-analysis of environmental contamination by phthalates, <http://www.ncbi.nlm.nih.gov/pubmed/23917738>, (2013)  
DOI: 10.1007/s11356-013-1982-5
14. Y Ji; F Wang; L Zhang; C Shan; Z Bai; Z Sun; L Liu; B Shen. A comprehensive assessment of human exposure to phthalates from environmental media and food in Tianjin, China. *J Hazard Mater* 279, 133-140 (2014)  
DOI: 10.1016/j.jhazmat.2014.06.055
15. RO Quintana-Belmares; AM Kraus; BK Esfahani; I Rosas-Pérez; D Mucs; R López-Marure; Å Bergman; E Alfaro-Moreno. Phthalate esters on urban airborne particles: Levels in PM10 and PM2.5 from Mexico City and theoretical assessment of lung exposure. *Environ Res* 161, 439-445 (2018)  
DOI: 10.1016/j.envres.2017.11.039
16. C Bergh; R Torgrip; G Emenius; C Östman. Organophosphate and phthalate esters in air and settled dust - a multi-location indoor study. *Indoor Air* 21, 67-76 (2011)  
DOI: 10.1111/j.1600-0668.2010.00684.x
17. H Fromme; T Lahrz; M Piloty; H Gebhart; A Oddoy; H Ruden. Occurrence of phthalates and musk fragrances in indoor air and dust from apartments and kindergartens in Berlin (Germany). *Indoor Air* 14, 188-195 (2004)  
DOI: 10.1111/j.1600-0668.2004.00223.x
18. E Diamanti-Kandarakis; JP Bourguignon; LC Giudice; R Hauser; GS Prins; AM Soto; RT Zoeller; AC Gore. Endocrine-disrupting chemicals: An Endocrine Society scientific statement. *Endocr Rev* 30, 293-342 (2009)  
DOI: 10.1210/er.2009-0002
19. I Colón; D Caro; CJ Bourdony; O Rosario. Identification of phthalate esters in the serum of young Puerto Rican girls with premature breast development. *Environ Health Perspect* 108, 895-900 (2000)  
DOI: 10.1289/ehp.00108895
20. SH Swan; KM Main; F Liu; SL Stewart; RL Kruse; AM Calafat; CS Mao; JB Redmon; CL Ternand; S Sullivan; JL Teague; Study for Future Families Research Team. Decrease in anogenital distance among male infants with prenatal phthalate exposure. *Environ Health Perspect* 113, 1056-1061 (2005)  
DOI: 10.1289/ehp.8100
21. SH Swan. Environmental phthalate exposure in relation to reproductive outcomes and other health endpoints in humans. *Environ Res* 108, 177-184 (2008)  
DOI: 10.1016/j.envres.2008.08.007
22. UN Joensen; H Frederiksen; MB Jensen; MP Lauritsen; IA Olesen; TH Lassen; A-M Andersson; N Jørgensen. Phthalate Excretion Pattern and Testicular Function: A Study of 881 Healthy Danish Men. *Environ Health Perspect* 120, 1397-1403 (2012)  
DOI: 10.1289/ehp.1205113
23. H Frederiksen; NE Skakkebaek; AM Andersson. Metabolism of phthalates in humans. *Mol Nutr Food Res* 51, 899-911 (2007)  
DOI: 10.1002/mnfr.200600243

## Phthalates as immunomodulators

24. SJ Kwack; EY Han; JS Park; JY Bae; IY Ahn; SK Lim; DH Kim; DE Jang; L Choi; HJ Lim; TH Kim; N Patra; KL Park; HS Kim; BM Lee. Comparison of the short-term toxicity of phthalate diesters and monoesters in sprague-dawley male rats. *Toxicol Res* 26, 75-82 (2010)  
DOI: 10.5487/TR.2010.26.1.075
25. T Okubo; T Suzuki; Y Yokoyama; K Kano; I Kano. Estimation of estrogenic and anti-estrogenic activities of some phthalate diesters and monoesters by MCF-7 cell proliferation assay *in vitro*. *Biol Pharm Bull* 26, 1219-1224 (2003)  
DOI: 10.1248/bpb.26.1219
26. H Zhang; Z Zhang; T Nakanishi; Y Wan; Y Hiromori; H Nagase; J Hu. Structure-Dependent Activity of Phthalate Esters and Phthalate Monoesters Binding to Human Constitutive Androstane Receptor. *Chem Res Toxicol* 28, 1196-1204 (2015)  
DOI: 10.1021/acs.chemrestox.5b00028
27. MT Bility; JT Thompson; RH McKee; RM David; JH Butala; JP Vanden Heuvel; JM Peters. Activation of mouse and human peroxisome proliferator-activated receptors (PPARs) by phthalate monoesters. *Toxicol Sci* 82, 170-182 (2004)  
DOI: 10.1093/toxsci/kfh253
28. N Heldring; A Pike; S Andersson; J Matthews; G Cheng; J Hartman; M Tujague; A Ström; E Treuter; M Warner; J-Å Gustafsson. Estrogen Receptors: How Do They Signal and What Are Their Targets. *Physiol Rev* 87, 905-931 (2007)  
DOI: 10.1152/physrev.00026.2006
29. J Matthews; JA Gustafsson. Estrogen signaling: a subtle balance between ER alpha and ER beta. *Mol Interv* 3, 281-292 (2003)  
DOI: 10.1124/mi.3.5.281
30. P Vrtačnik; B Ostanek; S Mencej-Bedrač; J Marc. The many faces of estrogen signaling. *Biochem medica* 24, 329-342 (2014)  
DOI: 10.11613/BM.2014.035
31. ER Levin. Membrane estrogen receptors signal to determine transcription factor function. *Steroids* 132, 1-4 (2018)  
DOI: 10.1016/j.steroids.2017.10.014
32. G Vail; TA Roepke. Membrane-initiated estrogen signaling via Gq-coupled GPCR in the central nervous system. *Steroids* 142, 77-83 (2019)  
DOI: 10.1016/j.steroids.2018.01.010
33. RA Davey; M Grossmann. Androgen Receptor Structure, Function and Biology: From Bench to Bedside. *Clin Biochem Rev* 37, 3-15 (2016)
34. NC Bennett; RA Gardiner; JD Hooper; DW Johnson; GC Gobe. Molecular cell biology of androgen receptor signalling. *Int J Biochem Cell Biol* 42, 813-827 (2010)  
DOI: 10.1016/j.biocel.2009.11.013
35. P Tontonoz; BM Spiegelman. Fat and Beyond: The Diverse Biology of PPAR $\gamma$ . *Annu Rev Biochem* 77, 289-312 (2008)  
DOI: 10.1146/annurev.biochem.-77.061307.091829
36. L la C Poulsen; M Siersbæk; S Mandrup. PPARs: Fatty acid sensors controlling metabolism. *Semin Cell Dev Biol* 23, 631-639 (2012)  
DOI: 10.1016/j.semcdb.2012.01.003
37. M Ahmadian; JM Suh; N Hah; C Liddle; AR Atkins; M Downes; RM Evans. Ppar $\gamma$  signaling and metabolism: The good, the

- bad and the future. *Nat Med* 19, 557-566 (2013)  
DOI: 10.1038/nm.3159
38. A Ammazalorso; C Maccallini; P Amoia; R Amoroso. Multitarget PPAR $\gamma$  agonists as innovative modulators of the metabolic syndrome. *Eur J Med Chem* 173, 261-273 (2019)  
DOI: 10.1016/j.ejmech.2019.04.030
39. X Ma; D Wang; W Zhao; L Xu. Deciphering the roles of PPAR $\gamma$  in adipocytes via dynamic change of transcription complex. *Front Endocrinol* 9, 473 (2018)  
DOI: 10.3389/fendo.2018.00473
40. L Larigot; L Juricek; J Dairou; X Coumoul. AhR signaling pathways and regulatory functions. *Biochim Open* 7, 1-9 (2018)  
DOI: 10.1016/j.biopen.2018.05.001
41. B Stockinger; P Di Meglio; M Gialitakis; JH Duarte. The Aryl Hydrocarbon Receptor: Multitasking in the Immune System. *Annu Rev Immunol* 32, 403-432 (2014)  
DOI: 10.1146/annurev-immunol-032713-120245
42. S Kovats. Estrogen receptors regulate innate immune cells and signaling pathways. *Cell Immunol* 294, 63-69 (2015)  
DOI: 10.1016/j.cellimm.2015.01.018
43. KL Phiel; RA Henderson; SJ Adelman; MM Elloso. Differential estrogen receptor gene expression in human peripheral blood mononuclear cell populations. *Immunol Lett* 97, 107-113 (2005)  
DOI: 10.1016/j.imlet.2004.10.007
44. D Khan; S Ansar Ahmed. The Immune System Is a Natural Target for Estrogen Action: Opposing Effects of Estrogen in Two Prototypical Autoimmune Diseases. *Front Immunol* 6, 635 (2015)  
DOI: 10.3389/fimmu.2015.00635
45. MC Erlandsson; C Ohlsson; JÅ Gustafsson; H Carlsten. Role of oestrogen receptors  $\alpha$  and  $\beta$  in immune organ development and in oestrogen-mediated effects on thymus. *Immunology* 103, 17-25 (2001)  
DOI: 10.1046/j.1365-2567.2001.01212.x
46. RJM Engler; MR Nelson; MM Klote; MJ VanRaden; CY Huang; NJ Cox; A Klimov; WA Keitel; KL Nichol; WW Carr; JJ Treanor. Half- vs full-dose trivalent inactivated influenza vaccine (2004-2005): Age, dose, and sex effects on immune responses. *Arch Intern Med* 168, 2405-2414 (2008)  
DOI: 10.1001/archinternmed.2008.513
47. MR Gubbels Bupp; TN Jorgensen. Androgen-Induced Immunosuppression. *Front Immunol* 9, 794 (2018)  
DOI: 10.3389/fimmu.2018.00794
48. JJ Lai; KP Lai; W Zeng; KH Chuang; S Altuwajiri; C Chang. Androgen receptor influences on body defense system via modulation of innate and adaptive immune systems: Lessons from conditional AR knockout mice. *Am J Pathol* 181, 1504-1512 (2012)  
DOI: 10.1016/j.ajpath.2012.07.008
49. TAP De Paula Couto; N Falsarella; CC De De Brandão Mattos; LC De Mattos. Total ige plasma levels vary according to gender and age in brazilian patients with allergic rhinitis. *Clinics* 69, 740-744 (2014)  
DOI: 10.6061/clinics/2014(11)06
50. L Széles; D Töröcsik; L Nagy. PPAR $\gamma$  in

- immunity and inflammation: cell types and diseases, *Biochim. Biophys. Acta, Mol. Basis Dis* 1771, 1014-1030 (2007)  
DOI: 10.1016/j.bbaliip.2007.02.005
51. M Ricote; AC Li; TM Willson; CJ Kelly; CK Glass. The peroxisome proliferator-activated receptor- $\gamma$  is a negative regulator of macrophage activation. *Nature* 391, 79-82 (1998)  
DOI: 10.1038/34178
  52. MA Bouhlel; B Derudas; E Rigamonti; R Dièvert; J Brozek; S Haulon; C Zawadzki; B Jude; G Torpier; N Marx; B Staels; G Chinetti-Gbaguidi. PPAR $\gamma$  Activation Primes Human Monocytes into Alternative M2 Macrophages with Anti-inflammatory Properties. *Cell Metab* 6, 137-143 (2007)  
DOI: 10.1016/j.cmet.2007.06.010
  53. SR Kim; KS Lee; HS Park; SJ Park; KH Min; SM Jin; YC Lee. Involvement of IL-10 in Peroxisome Proliferator-Activated Receptor  $\gamma$ -Mediated Anti-Inflammatory Response in Asthma. *Mol Pharmacol* 68, 1568-1575 (2005)  
DOI: 10.1124/mol.105.017160
  54. Y Kobayashi; S Ueki; G Mahemuti; T Chiba; H Oyamada; N Saito; A Kanda; H Kayaba; J Chihara. Physiological Levels of 15-Deoxy- $\Delta$  12,14 -Prostaglandin J 2 Prime Eotaxin-Induced Chemotaxis on Human Eosinophils through Peroxisome Proliferator-Activated Receptor- $\gamma$  Ligation. *J Immunol* 175, 5744-5750 (2005)  
DOI: 10.4049/jimmunol.175.9.5744
  55. SG Smith; H Imaoka; N Punia; A Irshad; LL Janssen; R Sehmi; GM Gauvreau. The Effect of PPAR Agonists on the Migration of Mature and Immature Eosinophils. *PPAR Res* 2012, 1-8 (2012)  
DOI: 10.1155/2012/235231
  56. SP Nobs; S Natali; L Pohlmeier; K Okreglicka; C Schneider; M Kurrer; F Sallusto; M Kopf. PPAR $\gamma$  in dendritic cells and T cells drives pathogenic type-2 effector responses in lung inflammation. *J Exp Med* 214, 3015-3035 (2017)  
DOI: 10.1084/jem.20162069
  57. H Hammad; HJ de Heer; T Soullié; V Angeli; F Trottein; HC Hoogsteden; BN Lambrecht. Activation of Peroxisome Proliferator-Activated Receptor- $\gamma$  in Dendritic Cells Inhibits the Development of Eosinophilic Airway Inflammation in a Mouse Model of Asthma. *Am J Pathol* 164, 263-271 (2004)  
DOI: 10.1016/S0002-9440(10)63116-1
  58. H Tuna; RG Avdiushko; VJ Sindhava; L Wedlund; CS Kaetzel; AM Kaplan; S Bondada; DA Cohen. Regulation of the mucosal phenotype in dendritic cells by PPAR $\gamma$ : role of tissue microenvironment. *J Leukoc Biol* 95, 471-485 (2014)  
DOI: 10.1189/jlb.0713408
  59. RB Clark; D Bishop-Bailey; T Estrada-Hernandez; T Hla; L Puddington; SJ Padula. The Nuclear Receptor PPAR $\gamma$  and Immunoregulation: PPAR $\gamma$  Mediates Inhibition of Helper T Cell Responses. *J Immunol* 164, 1364-1371 (2000)  
DOI: 10.4049/jimmunol.164.3.1364
  60. J Padilla; E Leung; RP Phipps. Human B lymphocytes and B lymphomas express PPAR- $\gamma$  and are killed by PPAR- $\gamma$  agonists. *Clin Immunol* 103, 22-33 (2002)  
DOI: 10.1006/clim.2001.5181
  61. C Gutiérrez-Vázquez; FJ Quintana. Regulation of the Immune Response by the Aryl Hydrocarbon Receptor. *Immunity* 48, 19-33 (2018)  
DOI: 10.1016/j.immuni.2017.12.012

## Phthalates as immunomodulators

62. A Kimura; T Naka; K Nohara; Y Fujii-Kuriyama; T Kishimoto. Aryl hydrocarbon receptor regulates Stat1 activation and participates in the development of Th17 cells. *Proc Natl Acad Sci U S A* 105, 9721-9726 (2008)  
DOI: 10.1073/pnas.0804231105
63. M Veldhoen; K Hirota; AM Westendorf; J Buer; L Dumoutier; JC Renaud; B Stockinger. The aryl hydrocarbon receptor links TH17-cell-mediated autoimmunity to environmental toxins. *Nature* 453, 106-109 (2008)  
DOI: 10.1038/nature06881
64. KN De Abrew; NE Kaminski; RS Thomas. An integrated genomic analysis of aryl hydrocarbon receptor-mediated inhibition of B-cell differentiation. *Toxicol Sci* 118, 454-469 (2010)  
DOI: 10.1093/toxsci/kfq265
65. FJ Quintana; DH Sherr. Aryl hydrocarbon receptor control of adaptive immunity *Pharmacol Rev* 65, 1148-1161 (2013)  
DOI: 10.1124/pr.113.007823
66. S Harris; SP Shubin; S Wegner; K Van Ness; F Green; SW Hong; EM Faustman. The presence of macrophages and inflammatory responses in an *in vitro* testicular co-culture model of male reproductive development enhance relevance to *in vivo* conditions. *Toxicol Vitro* 36, 210-215 (2016)  
DOI: 10.1016/j.tiv.2016.08.003
67. JF Hansen; K Bendtzen; M Boas; H Frederiksen; CH Nielsen; ÅK Rasmussen; U Feldt-Rasmussen. Influence of phthalates on cytokine production in monocytes and macrophages: a systematic review of experimental trials. *PLoS One* 10, e0120083 (2015)  
DOI: 10.1371/journal.pone.0120083
68. MB Bally; DJ Opheim; HG Shertzer. Di-(2-ethylhexyl) phthalate enhances the release of lysosomal enzymes from alveolar macrophages during phagocytosis. *Toxicology* 18, 49-60 (1980)  
DOI: 10.1016/0300-483X(80)90037-2
69. HG Shertzer; MB Bally; DJ Opheim. Inhibition of alveolar macrophage killing by di(2-ethylhexyl) phthalate. *Arch Environ Contam Toxicol* 14, 605-608 (1985)  
DOI: 10.1007/BF01055391
70. T Ohnishi; T Yoshida; A Igarashi; M Muroi; KI Tanamoto. Effects of possible endocrine disruptors on MyD88-independent TLR4 signaling. *FEMS Immunol Med Microbiol* 52, 293-295 (2008)  
DOI: 10.1111/j.1574-695X.2007.00355.x
71. C-C Hong; M Shimomura-Shimizu; M Muroi; K Tanamoto. Effect of endocrine disrupting chemicals on lipopolysaccharide-induced tumor necrosis factor-alpha and nitric oxide production by mouse macrophages. *Biol Pharm Bull* 27, 1136-1139 (2004)  
DOI: 10.1248/bpb.27.1136
72. E Koike; K Ichiro Inoue; R Yanagisawa; H Takano. Di-(2-ethylhexyl) phthalate affects immune cells from atopic prone mice *in vitro*. *Toxicology* 259, 54-60 (2009)  
DOI: 10.1016/j.tox.2009.02.002
73. X Pei; Z Duan; M Ma; Y Zhang; L Guo. Role of Ca/CaN/NFAT signaling in IL-4 expression by splenic lymphocytes exposed to phthalate (2-ethylhexyl) ester in spleen lymphocytes. *Mol Biol Rep* 41,

## Phthalates as immunomodulators

- 2129-2142 (2014)  
DOI: 10.1007/s11033-014-3062-4
74. P-S Oh; K-T Lim. Cudrania tricuspidata Bureau (CTB) Glycoprotein Inhibits Proliferation by Di(2-ethylhexyl) phthalate in Primary Splenocytes: Responses in Cell Proliferation Signaling. *Immunol Invest* 40, 339-355 (2011)  
DOI: 10.3109/08820139.2010.546468
75. U Yamashita; T Sugiura; Y Yoshida; E Kuroda. Effect of endocrine disrupters on thymocytes *in vitro*. *J UOEH* 25, 161-170 (2003)  
DOI: 10.7888/juoeh.25.161
76. JJ Schlezinger; GJ Howard; CH Hurst; JK Emberley; DJ Waxman; T Webster; DH Sherr. Environmental and Endogenous Peroxisome Proliferator-Activated Receptor  $\gamma$  Agonists Induce Bone Marrow B Cell Growth Arrest and Apoptosis: Interactions between Mono(2-ethylhexyl) phthalate, 9- cis -Retinoic Acid, and 15-Deoxy- $\Delta$  12,14 -prostaglandin J 2 . *J Immunol* 173, 3165-3177 (2004)  
DOI: 10.4049/jimmunol.173.5.3165
77. K Martins; B Applegate; B Hagedorn; J Kennish; P Zwollo. Di(2-ethylhexyl) phthalate inhibits B cell proliferation and reduces the abundance of IgM-secreting cells in cultured immune tissues of the rainbow trout. *Fish Shellfish Immunol* 44, 332-341 (2015)  
DOI: 10.1016/j.fsi.2015.02.037
78. R Nakamura; R Teshima; J ichi Sawada. Effect of dialkyl phthalates on the degranulation and Ca<sup>2+</sup> response of RBL-2H3 mast cells. *Immunol Lett* 80, 119-124 (2002)  
DOI: 10.1016/S0165-2478(01)00318-2
79. C Glue; A Millner; U Bodtger; T Jinquan; LK Poulsen. *In vitro* effects of monophthalates on cytokine expression in the monocytic cell line THP-1 and in peripheral blood mononuclear cells from allergic and non-allergic donors. *Toxicol In vitro* 16, 657-662 (2002)  
DOI: 10.1016/S0887-2333(02)00082-6
80. HG Kim; S min Yeon; KH Kim; H Kim; J Il Park; HJ Kang; EJ Cha; HD Park; HJ Kang; TW Park; YH Jeon; Y In Park; KT Chang; YW Jung. Estrogenic Endocrine-Disrupting Chemicals Modulate the Production of Inflammatory Mediators and Cell Viability of Lipopolysaccharide-Stimulated Macrophages. *Inflammation* 38, 595-605 (2015)  
DOI: 10.1007/s10753-014-9966-2
81. KH Kim; SM Yeon; HG Kim; HS Choi; H Kang; HD Park; TW Park; SP Pack; EH Lee; Y Byun; SE Choi; KS Lee; UH Ha; YW Jung. Diverse influences of androgen-disrupting chemicals on immune responses mounted by macrophages. *Inflammation* 37, 649-656 (2014)  
DOI: 10.1007/s10753-013-9781-1
82. L Li; HS Li; NN Song; HM Chen. The immunotoxicity of dibutyl phthalate on the macrophages in mice. *Immunopharmacol Immunotoxicol* 35, 272-281 (2013)  
DOI: 10.3109/08923973.2013.768267
83. A Kocbach Bølling; J Ovrevik; JT Samuelsen; JA Holme; KE Rakkestad; GH Mathisen; RE Paulsen; M Suárez Korsnes; R Becher. Mono-2-ethylhexylphthalate (MEHP) induces TNF- $\alpha$  release and macrophage differentiation through different signalling pathways in RAW264.7 cells. *Toxicol Lett* 209, 43-50 (2012)  
DOI: 10.1016/j.toxlet.2011.11.016

## Phthalates as immunomodulators

84. T Ito; KI Inoue; N Nishimura; H Takano. Phthalate esters modulate the differentiation and maturation of mouse peripheral blood mononuclear cell-derived dendritic cells. *J Appl Toxicol* 32, 142-148 (2012)  
DOI: 10.1002/jat.1652
85. KE Rakkestad; JA Holme; RE Paulsen; PE Schwarze; R Becher. Mono(2-ethylhexyl) phthalate induces both pro- and anti-inflammatory responses in rat alveolar macrophages through crosstalk between p38, the lipoxygenase pathway and PPAR $\alpha$ . *Inhal Toxicol* 22, 140-150 (2010)  
DOI: 10.3109/08958370903019885
86. J Naarala; A Korpi. Cell death and production of reactive oxygen species by murine macrophages after short term exposure to phthalates. *Toxicol Lett* 188, 157-160 (2009)  
DOI: 10.1016/j.toxlet.2009.04.001
87. MH Lee; J Park; SW Chung; BY Kang; SH Kim; TS Kim. Enhancement of interleukin-4 production in activated CD4+ T cells by diphthalate plasticizers via increased NF-AT binding activity. *Int Arch Allergy Immunol* 134, 213-222 (2004)  
DOI: 10.1159/000078768
88. T Gourlay; I Samartzis; D Stefanou; K Taylor. Inflammatory response of rat and human neutrophils exposed to Di-(2-ethyl-hexyl)-phthalate-plasticized polyvinyl chloride. *Artificial Organs* 27, 256-260 (2003)  
DOI: 10.1046/j.1525-1594.2003.07107.x
89. H Watanuki; Y Gushiken; M Sakai. *In vitro* modulation of common carp (*Cyprinus carpio* L.) phagocytic cells by Di-n-butyl phthalate and Di-2-ethylhexyl phthalate. *Aquat Toxicol* 63, 119-126 (2003)  
DOI: 10.1016/S0166-445X(02)00172-8
90. IA Tavares; ND Vine. Phthalic acid esters inhibit arachidonate metabolism by rat peritoneal leucocytes. *J Pharm Pharmacol* 37, 67-68 (1985)  
DOI: 10.1111/j.2042-7158.1985.tb04936.x
91. A Bennasroune; L Rojas; L Foucaud; S Goulaouic; P Laval-Gilly; M Fickova; N Couleau; C Durandet; S Henry; J Falla. Effects of 4-nonylphenol and/or diisononylphthalate on THP-1 cells: Impact of endocrine disruptors on human immune system parameters. *Int J Immunopathol Pharmacol* 25, 365-376 (2012)  
DOI: 10.1177/039463201202500206
92. N Couleau; J Falla; A Beillerot; E Battaglia; M D'Innocenzo; S Plançon; P Laval-Gilly; A Bennasroune. Effects of Endocrine Disruptor Compounds, Alone or in Combination, on Human Macrophage-Like THP-1 Cell Response. *PLoS One* 10, e0131428 (2015)  
DOI: 10.1371/journal.pone.0131428
93. LM Tetz; DM Aronoff; R Loch-Carus. Mono-ethylhexyl phthalate stimulates prostaglandin secretion in human placental macrophages and THP-1 cells. *Reprod Biol Endocrinol* 13, 56 (2015)  
DOI: 10.1186/s12958-015-0046-8
94. D Teixeira; C Marques; D Pestana; A Faria; S Norberto; C Calhau; R Monteiro. Effects of xenoestrogens in human M1 and M2 macrophage migration, cytokine release, and estrogen-related signaling pathways. *Environ Toxicol* 31, 1496-1509 (2016)  
DOI: 10.1002/tox.22154

## Phthalates as immunomodulators

95. M Miyamoto; S Sasakawa. Effects of Plasticizers and Plastic Bags on Granulocyte Function during Storage. *Vox Sang* 53, 19-22 (1987)  
DOI: 10.1111/j.1423-0410.1987.tb04907.x
96. S Palleschi; B Rossi; L Diana; L Silvestroni. Di(2-ethylhexyl) phthalate stimulates Ca<sup>2+</sup> entry, chemotaxis and ROS production in human granulocytes. *Toxicol Lett* 187, 52-57 (2009)  
DOI: 10.1016/j.toxlet.2009.01.031
97. CA Rosado-Berrios; C Vélez; B Zayas. Mitochondrial permeability and toxicity of diethylhexyl and monoethylhexyl phthalates on TK6 human lymphoblasts cells. *Toxicol Vitr* 25, 2010-2016 (2011)  
DOI: 10.1016/j.tiv.2011.08.001
98. G Sager; C Little. The effect of the plasticizers TBEP (tris-(2-butoxyethyl)-phosphate) and DHEP (DI-(2-ethylhexyl)-phthalate) on  $\beta$ -adrenergic ligand binding to  $\alpha$ 1-acid glycoprotein and mononuclear leukocytes. *Biochem Pharmacol* 38, 2551-2557 (1989)  
DOI: 10.1016/0006-2952(89)90101-9
99. NH Kleinsasser; BC Wallner; ER Kastenbauer; H Weissacher; UA Harreus. Genotoxicity of di-butyl-phthalate and di-iso-butyl-phthalate in human lymphocytes and mucosal cells. *Teratog Carcinog Mutagen* 21, 189-196 (2001)  
DOI: 10.1002/tcm.1007
100. NH Kleinsasser; UA Harréus; ER Kastenbauer; BC Wallner; AW Sassen; R Staudenmaier; AW Rettenmeier. Mono(2-ethylhexyl)phthalate exhibits genotoxic effects in human lymphocytes and mucosal cells of the upper aerodigestive tract in the comet assay. *Toxicol Lett* 148, 83-90 (2004)  
DOI: 10.1016/j.toxlet.2003.12.013
101. AM Vetrano; DL Laskin; F Archer; K Syed; JP Gray; JD Laskin; N Nwebube; B Weinberger. Inflammatory effects of phthalates in neonatal neutrophils. *Pediatr Res* 68, 134-139 (2010)  
DOI: 10.1203/PDR.0b013e3181e5c1f7
102. J Lee; PS Oh; KT Lim. Allergy-related cytokines (IL-4 and TNF- $\alpha$ ) are induced by Di(2-ethylhexyl) phthalate and attenuated by plant-originated glycoprotein (75 kDa) in HMC-1 cells. *Environ Toxicol* 26, 364-372 (2011)  
DOI: 10.1002/tox.20563
103. J Nishioka; C Iwahara; M Kawasaki; F Yoshizaki; H Nakayama; K Takamori; H Ogawa; K Iwabuchi. Di-(2-ethylhexyl) phthalate induces production of inflammatory molecules in human macrophages. *Inflamm Res* 61, 69-78 (2012)  
DOI: 10.1007/s00011-011-0390-x
104. CH Kuo; CC Hsieh; HF Kuo; MY Huang; SN Yang; LC Chen; SK Huang; CH Hung. Phthalates suppress type I interferon in human plasmacytoid dendritic cells via epigenetic regulation. *Allergy Eur J Allergy Clin Immunol* 68, 870-879 (2013)  
DOI: 10.1111/all.12162
105. YL Hsu; JY Hung; EM Tsai; CY Wu; YW Ho; SF Jian; MC Yen; WA Chang; MF Hou; PL Kuo. Benzyl butyl phthalate increases the chemoresistance to doxorubicin/cyclophosphamide by increasing breast cancer-associated dendritic cell-derived CXCL1/GRO $\alpha$  and S100A8/A9. *Oncol Rep* 34, 2889-2900 (2015)  
DOI: 10.3892/or.2015.4307

## Phthalates as immunomodulators

106. Y Han; X Wang; G Chen; G Xu; X Liu; W Zhu; P Hu; Y Zhang; C Zhu; J Miao. Di-(2-ethylhexyl) phthalate adjuvantly induces imbalanced humoral immunity in ovalbumin-sensitized BALB/c mice ascribing to T follicular helper cells hyperfunction. *Toxicology* 324, 88-97 (2014)  
DOI: 10.1016/j.tox.2014.07.011
107. ST Larsen; GD Nielsen. The adjuvant effect of di-(2-ethylhexyl) phthalate is mediated through a PPAR $\alpha$ -independent mechanism. *Toxicol Lett* 170, 223-228 (2007)  
DOI: 10.1016/j.toxlet.2007.03.009
108. ST Larsen; JS Hansen; EW Hansen; PA Clausen; GD Nielsen. Airway inflammation and adjuvant effect after repeated airborne exposures to di-(2-ethylhexyl) phthalate and ovalbumin in BALB/c mice. *Toxicology* 235, 119-129 (2007)  
DOI: 10.1016/j.tox.2007.03.010
109. JS Hansen; ST Larsen; LK Poulsen; GD Nielsen. Adjuvant effects of inhaled mono-2-ethylhexyl phthalate in BALB/cJ mice. *Toxicology* 232, 79-88 (2007)  
DOI: 10.1016/j.tox.2006.12.011
110. YH Hwang; MJ Paik; ST Yee. Diisononyl phthalate induces asthma via modulation of Th1/Th2 equilibrium. *Toxicol Lett* 272, 49-59 (2017)  
DOI: 10.1016/j.toxlet.2017.03.014
111. J Guo; B Han; L Qin; B Li; H You; J Yang; D Liu; C Wei; E Nanberg; CG Bornehag; X Yang. Pulmonary toxicity and adjuvant effect of Di-(2-ethylhexyl) phthalate in Ovalbumin-immunized BALB/c mice. *PLoS One* 7, e39008 (2012)  
DOI: 10.1371/journal.pone.0039008
112. K Martins; B Hagedorn; S Ali; J Kennish; B Applegate; M Leu; L Epp; C Pallister; P Zwollo. Tissue Phthalate Levels Correlate With Changes in Immune Gene Expression in a Population of Juvenile Wild Salmon. *Arch Environ Contam Toxicol* 71, 35-47 (2016)  
DOI: 10.1007/s00244-016-0283-7
113. J Li; L Li; H Zuo; C Ke; B Yan; H Wen; Y Zhang; X Yang. T-helper type-2 contact hypersensitivity of balb/c mice aggravated by dibutyl phthalate via long-term dermal exposure. *PLoS One* 9, e87887 (2014)  
DOI: 10.1371/journal.pone.0087887
114. M He; KI Inoue; S Yoshida; M Tanaka; H Takano; G Sun; T Ichinose. Effects of airway exposure to di-(2-ethylhexyl) phthalate on allergic rhinitis. *Immunopharmacol Immunotoxicol* 35, 390-395 (2013)  
DOI: 10.3109/08923973.2013.787432
115. E Koike; R Yanagisawa; K Sadakane; KI Inoue; T Ichinose; H Takano. Effects of diisononyl phthalate on atopic dermatitis *in vivo* and immunologic responses *in vitro*. *Environ Health Perspect* 118, 472-478 (2010)  
DOI: 10.1289/ehp.0901255
116. T Maruyama; T Shiba; H Iizuka; T Matsuda; K Kurohane; Y Imai. Effects of phthalate esters on dendritic cell subsets and interleukin-4 production in fluorescein isothiocyanate-induced contact hypersensitivity. *Microbiol Immunol* 51, 321-326 (2007)  
DOI: 10.1111/j.1348-0421.2007.tb03914.x
117. T Shigeno; M Katakuse; T Fujita; Y Mukoyama; H Watanabe. Phthalate ester-induced thymic stromal

## Phthalates as immunomodulators

- lymphopoietin mediates allergic dermatitis in mice. *Immunology* 128 (1 Suppl), e849-e857 (2009)  
DOI: 10.1111/j.1365-2567.2009.03094.x
118. RP Larson; SC Zimmerli; MR Comeau; A Itano; M Omori; M Iseki; C Hauser; SF Ziegler. Dibutyl Phthalate-Induced Thymic Stromal Lymphopoietin Is Required for Th2 Contact Hypersensitivity Responses. *J Immunol* 184, 2974-2984 (2010)  
DOI: 10.4049/jimmunol.0803478
119. Y Imai; A Kondo; H Iizuka; T Maruyama; KK Kurohane. Effects of phthalate esters on the sensitization phase of contact hypersensitivity induced by fluorescein isothiocyanate. *Clin Exp Allergy* 36, 1462-1468 (2006)  
DOI: 10.1111/j.1365-2222.2006.02574.x
120. JF Zhao; SH Hsiao; MH Hsu; KC Pao; YR Kou; SK Shyue; TS Lee. Di-(2-ethylhexyl) phthalate accelerates atherosclerosis in apolipoprotein E-deficient mice. *Arch Toxicol* 90, 181-190 (2016)  
DOI: 10.1007/s00204-014-1377-5
121. CJ Murphy; AR Stermer; JH Richburg. Age- and Species-Dependent Infiltration of Macrophages into the Testis of Rats and Mice Exposed to Mono-(2-Ethylhexyl) Phthalate (MEHP)1. *Biol Reprod* 91, 18 (2014)  
DOI: 10.1095/biolreprod.113.115527
122. M Kitaoka; S Hirai; H Terayama; M Naito; N Qu; N Hatayama; H Miyaso; Y Matsuno; M Komiyama; M Itoh; C Mori. Effects on the local immunity in the testis by exposure to di-(2-ethylhexyl) phthalate (DEHP) in mice. *J Reprod Dev* 59, 485-490 (2013)  
DOI: 10.1262/jrd.2012-180
123. SJ Zheng; HJ Tian; J Cao; YQ Gao. Exposure to di(n-butyl)phthalate and benzo(a)pyrene alters IL-1 $\beta$  secretion and subset expression of testicular macrophages, resulting in decreased testosterone production in rats. *Toxicol Appl Pharmacol* 248, 28-37 (2010)  
DOI: 10.1016/j.taap.2010.07.008
124. S Hirai; M Naito; M Kuramasu; Y Ogawa; H Terayama; N Qu; N Hatayama; S Hayashi; M Itoh. Low-dose exposure to di-(2-ethylhexyl) phthalate (DEHP) increases susceptibility to testicular autoimmunity in mice. *Reprod Biol* 15, 163-171 (2015)  
DOI: 10.1016/j.repbio.2015.06.004
125. LM Connor; SC Tang; E Cognard; S Ochiai; KL Hilligan; SI Old; C Pellefigues; RF White; D Patel; AAT Smith; DA Eccles; O Lamiabile; MJ McConnell; F Ronchese. Th2 responses are primed by skin dendritic cells with distinct transcriptional profiles. *J Exp Med* 214, 125-142 (2017)  
DOI: 10.1084/jem.20160470
126. H Xu; X Dong; Z Zhang; M Yang; X Wu; H Liu; Q Lao; C Li. Assessment of immunotoxicity of dibutyl phthalate using live zebrafish embryos. *Fish Shellfish Immunol* 45, 286-292 (2015)  
DOI: 10.1016/j.fsi.2015.04.033
127. M Tanaka; KI Inoue; T Momoi; H Takano. *In vivo* immunoamplifying effects of di-(2-ethylhexyl) phthalate on cytokine response. *Immunopharmacol Immunotoxicol* 35, 147-150 (2013)  
DOI: 10.3109/08923973.2012.733705
128. ECM Tonk; A Verhoef; ER Gremmer; H van Loveren; AH Piersma. Relative sensitivity of developmental and immune parameters in juvenile versus adult male

## Phthalates as immunomodulators

- rats after exposure to di(2-ethylhexyl) phthalate. *Toxicol Appl Pharmacol* 260, 48-57 (2012)  
DOI: 10.1016/j.taap.2012.01.018
129. T Matsuda; K Kurohane; Y Imai. Di-(2-ethylhexyl) phthalate enhances skin sensitization to isocyanate haptens in mice. *Toxicol Lett* 192, 97-100 (2010)  
DOI: 10.1016/j.toxlet.2009.10.009
130. U Yamashita; E Kuroda; Y Yoshida; T Sugiura. Effect of Endocrine Disrupters on Immune Responses *In vivo*. *J UOEH* 25, 365-374 (2003)  
DOI: 10.7888/juoeh.25.365
131. I Rusyn; MB Kadiiska; A Dikalova; H Kono; M Yin; K Tsuchiya; RP Mason; JM Peters; FJ Gonzalez; BH Segal; SM Holland; RG Thurman. Phthalates rapidly increase production of reactive oxygen species *in vivo*: Role of Kupffer cells. *Mol Pharmacol* 59, 744-750 (2001)  
DOI: 10.1124/mol.59.4.744
132. RM Whyatt; MS Perzanowski; AC Just; AG Rundle; KM Donohue; AM Calafat; LA Hoepner; FP Perera; RL Miller. Asthma in Inner-City Children at 5-11 Years of Age and Prenatal Exposure to Phthalates: The Columbia Center for Children's Environmental Health Cohort. *Environ Health Perspect* 122, 1141-1146 (2014)  
DOI: 10.1289/ehp.1307670
133. M Gascon; M Casas; E Morales; D Valvi; A Ballesteros-Gómez; N Luque; S Rubio; N Monfort; R Ventura; D Martínez; J Sunyer; M Vrijheid. Prenatal exposure to bisphenol A and phthalates and childhood respiratory tract infections and allergy. *J Allergy Clin Immunol* 135, 370-378 (2015)  
DOI: 10.1016/j.jaci.2014.09.030
134. HY Ku; PH Su; HJ Wen; HL Sun; CJ Wang; HY Chen; JJK Jaakkola; S-L Wang; T Group. Prenatal and Postnatal Exposure to Phthalate Esters and Asthma: A 9-Year Follow-Up Study of a Taiwanese Birth Cohort. *PLoS One* 10, e0123309 (2015)  
DOI: 10.1371/journal.pone.0123309
135. MH Soomro; N Baiz; C Philippat; C Vernet; V Siroux; C Nichole Maesano; S Sanyal; R Slama; C-G Bornehag; I Annesi-Maesano. Prenatal Exposure to Phthalates and the Development of Eczema Phenotypes in Male Children: Results from the EDEN Mother-Child Cohort Study. *Environ Health Perspect* 126, 027002 (2018)  
DOI: 10.1289/EHP1829
136. J Ashley-Martin; L Dodds; AR Levy; RW Platt; JS Marshall; TE Arbuckle. Prenatal exposure to phthalates, bisphenol A and perfluoroalkyl substances and cord blood levels of IgE, TSLP and IL-33. *Environ Res* 140, 360-368 (2015)  
DOI: 10.1016/j.envres.2015.04.010
137. IJ Wang; CC Lin; YJ Lin; WS Hsieh; PC Chen. Early life phthalate exposure and atopic disorders in children: A prospective birth cohort study. *Environ Int* 62, 48-54 (2014)  
DOI: 10.1016/j.envint.2013.09.002
138. JA Hoppin; R Jaramillo; SJ London; RJ Bertelsen; PM Salo; DP Sandler; DC Zeldin. Phthalate exposure and allergy in the U.S. population: Results from NHANES 2005-2006. *Environ Health Perspect* 121, 1129-1134 (2013)  
DOI: 10.1289/ehp.1206211
139. T Deuschie; R Reiter; W Butte; B Heinzow; T Keck; H Riechelmann. A controlled challenge study on Di(2-

ethylhexyl) phthalate (DEHP) in house dust and the immune response in human nasal mucosa of allergic subjects. *Environ Health Perspect* 116, 1487-1493 (2008)  
DOI: 10.1289/ehp.11474

**Abbreviations:** EDCs: Endocrine disrupting compounds; PPAR-gamma: Peroxisome proliferator-activated receptor gamma; AhR: Aryl hydrocarbon receptor; MHC: Major histocompatibility complex; PVC: Polyvinyl chloride; ug: micrograms; Bw: body weight; PM2.5: Fine particulate matter (of less than 2.5 micrometers); DEHP: Bis-(2-ethylhexyl) phthalate; DBP: Di-n-butyl phthalate; EPA: Environmental Protection Agency; MEHP: Mono(2-ethylhexyl) phthalate; ER: Estrogen receptor; AR: Androgen receptor; EREs: Estrogen response elements; DNA: Deoxyribonucleic acid; eRs: Estrogen receptors; GPER: G protein receptor-coupled estrogen receptor; cAMP: Cyclic adenosine monophosphate; DAG: Diacylglycerol; IP3: Inositol triphosphate; DHT: Dihydrotestosterone; AREs: Androgen response elements; ERK: Extracellular signal-regulated kinase; MAPK: Mitogen-activated protein kinase; RXR: Retinoid X receptor; PPRE: PPAR response element; SMRT: silencing mediator of retinoid and thyroid hormone receptors; nCoR: nuclear receptor corepressor; HDACs: histone deacetylases; CBP: CREB-binding protein; PBP: PPAR-binding protein 4; TCDD: 2,3,7,8-tetrachlorodibenzo-p-dioxin; bHLH: basic helix-loop-helix; HSP90: heat-shock protein 90; AIP: Aryl hydrocarbon receptor-interacting protein; ARNT: Aryl hydrocarbon nuclear translocator; DRE: Dioxin response element; CYP1A1: Cytochrome P450, family 1, subfamily A, polypeptide 1; AhRR: AhR receptor; SW1/SNF: SWItch/Sucrose Non-Fermentable remodeling complex; SRC-1: steroid receptor coactivator-1; FAK: focal adhesion kinase; NK: natural killer

cells; Th: T helper cells; Treg: Regulatory T cells; HLA: human leukocyte antigen; G-CSF: Granulocyte colony stimulating factor; IL: interleukin; IgE: immunoglobulin E; 15-dPGJ<sub>2</sub>: 15-deoxy- $\Delta$ 12,14-prostaglandin J<sub>2</sub>; IFN-: Interferon; iNOS: Inducible Nitric oxide synthase; M1/M2: macrophage phenotype 1 or 2; CD: cluster of differentiation; BM-DC: Bone marrow-derived dendritic cells; iTregs: Induced Tregs; CCR9: C-C chemokine receptor type 9; mDC: mature dendritic cells; PMA: phorbol myristate acetate; TUNEL: Terminal deoxynucleotidyl transferase dUTP nick end labeling assay; DEP: Diethyl phthalate; TNF: tumor necrosis factor; CXCL8: C-X-C chemokine ligand 8; DnBP: Di-n-butyl phthalate; BzBP: Benzylbutyl phthalate; DPRP: Dipropyl phthalate; TCR: T cell receptor; Ca: calcium; CaN: calmodulin; NF-AT: Nuclear factor of activated T-cells; PKC: protein kinase C; NF-kappa B: nuclear factor kappa B; DiBP: Diisobutyl phthalate; PBMC: peripheral blood mononuclear cells; fMLP: N-formyl-methionyl-leucyl-phenylalanine; COX: cyclooxygenase; Thf: follicular T helper cells; DPP: D-n-propyl phthalate; DINP: Diisononyl phthalate; TSLP: Thymic stromal lymphopoietin; BBzP: Benzylbutyl phthalate; MBzP: Monobenzyl phthalate; DiNP: Diisononyl phthalate; MCPP: Mono (3-carboxypropyl) phthalate; MEP: Monoethyl phthalate; MBP: Monobutyl phthalate; ConA: concanavalin A; DIBP: Diisobutyl phthalate; cx-MiDP: Carboxi-monoisodecyl phthalate, DAP: Diallyl phthalate, DCHP: Dicyclohexyl phthalate, DHP: Di-n-hexyl phthalate, DiHepP: Diisoheptyl phthalate, DiHP: Diisohexyl phthalate, DiP: Diisononyl phthalate, DiPP: Diisopentyl phthalate, DnOP: Di-n-octyl phthalate, DnPP: Di-n-pentyl phthalate, DPeP: Di-n-pentyl phthalate, DTDP: Ditridecyl phthalate, MBEP: Monobenzyl phthalate, MBUP: Mono-n-butyl phthalate, MIDP : Monoisodecyl phthalate, MiDP: Monoisodecyl phthalate, MINP : Monoisononyl phthalate, MnBP: Mono-n-butyl phthalate, MOP : Mono-n-

## **Phthalates as immunomodulators**

octyl phthalate, MTDP: Monotridecyl phthalate, ROS: Reactive oxygen species, LTB<sub>4</sub>: leukotriene B<sub>4</sub>; PGE-2: prostaglandin 2; STAT: Signal transducer and activator of transcription, TBEP : Tris-(2-butoxyethyl)-phosphate

**Key Words:** Phthalates, Immunomodulation, Immune, Endocrine, Disruptor, EDC, Review

**Send correspondence to:** Karen Elizabeth Nava-Castro, Laboratorio de Genotoxicología y Mutagenesis Ambientales, Departamento de Ciencias Ambientales. Centro de Ciencias de la Atmosfera, Universidad Nacional Autónoma de México, CP 04510, Ciudad de México, México, Tel: 52-1-55-56224077, Fax: 52-1-55-56223369, E-mail: karlenc@atmosfera.unam.mx